

Mating balls in stoneflies (Insecta, Plecoptera)

Las bolas de apareamiento en plecópteros (Insecta, Plecoptera)

J.M. TIERNO DE FIGUEROA (1), J.M. LUZÓN-ORTEGA (2) & M.J. LÓPEZ-RODRÍGUEZ (1)

(1) Departamento de Biología Animal. Facultad de Ciencias. Universidad de Granada. 18071. Granada (España). E-mail: jmtdef@ugr.es, manujlr@ugr.es

(2) Hydraena S.L.L. C/ Nenúfares, 8. 18213, Jun (Granada, España).
E-mail: julioluzon@hydraena.com

Recibido el 29 de agosto de 2006. Aceptado el 15 de septiembre de 2006.

ISSN: 1130-4251 (2006), vol. 17, 93-96

Mating balls occur when some males simultaneously attempt copulation with one female and they pile onto and around her forming a ball. Mating balls have been described in different animal groups, particularly in colubrids and anurans (Madsen & Shine, 1993; Luiselli, 1996; Wells, 1977; Reyer *et al.*, 2003; Pleguezuelos & Feriche, 2006). As pointed out by Shine *et al.* (2000) for snakes, mating balls are most likely to occur in species that (1) do not display male-male combat, (2) have high population densities, and especially (3) have mating periods soon after spring emergence, while the snakes are highly aggregated at overwintering dens. Moreover, in all these animals females are larger than males.

All these criteria are shared by many Plecoptera species in our latitudes: no existence of male-male combat, high population density after the emergence (the emergence is usually very concentrated in a short period of time) and tendency to aggregation (Hanada *et al.*, 1994; Stewart, 1994; Alexander & Stewart, 1996; Jaímez-Cuéllar & Tierno de Figueroa, 2005).

Up to now, the attempts of one male trying to displace a mating male and mate guarding behaviours have been described for some different stonefly species (Alexander & Stewart, 1996; Tierno de Figueroa *et al.*, 2000; Tierno de Figueroa, 2003), but not forming a real mating ball.

Brinck (1949) pointed that when a female of a setipalpian species with short-winged males has emerged, she usually becomes surrounded by waiting males; this has been observed in *Diura bicaudata* (Linnaeus,

1758), *Arcynopteryx compacta* (McLachland, 1872) and *Dinocras cephalotes* (Curtis, 1827). Nevertheless, according with his description, the remaining males (not the one copulating) wait around the female sides (no attempts of copulation were observed until the male from the pair retired and left a place).

We have observed in the field the formation of mating balls in two different stonefly species: *Dinocras cephalotes* (observed at Río Peguera, Pyrenees, Lérida province, Spain, June, 24th 2006) (Fig. 1) involving five brachypterous males, and *Hemimelaena flaviventris* (Pictet, 1842) (observed in Río Despeñaperros, Sierra Morena, Jaén province, Spain, April, 27th 2006) (Fig. 2) involving three macropterus males. These data constitute the first evidence of this behaviour in Plecoptera. Probably, and in relation with the little research devoted to mate-finding, copulation and displacement attempts during copulation in stoneflies (Tierno de Figueroa, 2003),



Fig. 1.—Mating ball of *Dinocras cephalotes*. The female can be distinguished by its long wings, while males are all brachypterous.

Fig. 1.—Bola de apareamiento de *Dinocras cephalotes*. La hembra se puede distinguir por sus largas alas, mientras que los machos son braquípteros.



Fig. 2.—Mating ball of *Hemimelaena flaviventris*. The female can be distinguished by its higher size and the presence of the subgenital plate.

Fig. 2.—Bola de apareamiento de *Hemimelaena flaviventris*. La hembra se puede distinguir por su mayor tamaño y por la presencia de la placa subgenital.

the existence of mating balls in Plecoptera is probably more common than currently thought.

REFERENCES

- ALEXANDER, K.D. & STEWART, K.W. 1996. Description and theoretical considerations of mate-finding and other adult behaviours in a Colorado population of *Claassenia sabulosa* (Plecoptera, Perlidae). *Annals of the Entomological Society of America*, 89: 290-296.
- BRINCK, P. 1949. Studies on Swedish stoneflies (Plecoptera). *Opuscula Entomologica*, suppl. 11: 1-250.
- HANADA, S.; ISOBE, Y.; WADA, K. & NAGOSHI, M. 1994. Drumming behavior of two stonefly species, *Microperla brevicauda* (Peltoperlidae) and *Kamimuria tibialis* (Pictet) (Perlidae) in relation to other behaviors. *Aquatic Insects*, 16 (2): 75-89.
- JÁIMEZ-CUÉLLAR, P. & TIERNO DE FIGUEROA, J.M. 2005. Local emergence patterns in *Dinocras cephalotes* (Plecoptera, Perlidae). *Biologia, Bratislava*, 60 (5): 539-543.

- LUISELLI, L. 1996. Individual success in mating balls of the grass snake, *Natrix natrix*: size is important. *Journal of Zoology*, 239: 731-740.
- MADSEN, T. & SHINE, R. G. 1993. Male mating success and body size in European grass snake. *Copeia*, 1993: 561-564.
- PLEGUEZUELOS, J.M. & FERICHE, M. 2006. Reproductive ecology of a Mediterranean ratsnake, the Ladder snake *Rhinechis scalaris* (Schinz, 1822). *Herpetological Journal*, 16: 177-182.
- REYER, H.U.; NIEDERER, B. & HETTYEV, A. 2003. Variation in fertilization abilities between hemiclelonal hybrid and sexual parental males of sympatric water frogs (*Rana lessonae*, *R. esculenta*, *R. ridibunda*). *Behavioural Ecology and Sociobiology*, 54: 274-284.
- SHINE, R.; OLSSON, M.M.; MOORE, I.T.; LEMASTER, M.P.; GREENE, M. & MASON, R.T. 2000. Body size enhances mating success in male garter snakes. *Animal Behaviour*, 59: F4-F11.
- STEWART, K.W. 1994. Theoretical considerations of mate finding and other adult behaviours of Plecoptera. *Aquatic Insects*, 16: 95-104.
- TIERNOW DE FIGUEROA, J.M. 2003. Mate guarding and displacement attempts in stoneflies (Insecta, Plecoptera). *Biologia, Bratislava*, 58: 925-928.
- TIERNOW DE FIGUEROA, J.M.; LUZÓN-ORTEGA, J.M. & SÁNCHEZ-ORTEGA, A. 2000. Calling, mating and oviposition in *Isoperla curtata* (Plecoptera, Perlodidae). *European Journal of Entomology*, 97: 171-175.
- WELLS, K.D. 1977. The social behaviour of anuran amphibians. *Animal Behaviour*, 25: 666-693.