

RESEARCH
REVIEW



Reassessing global change research priorities in mediterranean terrestrial ecosystems: how far have we come and where do we go from here?

E. Doblas-Miranda^{1*}, J. Martínez-Vilalta^{1,2}, F. Lloret^{1,2}, A. Álvarez¹, A. Ávila¹, F. J. Bonet³, L. Brotons^{1,4}, J. Castro⁵, J. Curiel Yuste⁶, M. Díaz⁶, P. Ferrandis⁷, E. García-Hurtado⁸, J. M. Iriondo⁹, T. F. Keenan¹⁰, J. Latron¹¹, J. Llusà^{1,12}, L. Loepfe¹³, M. Mayol¹, G. Moré¹, D. Moya⁷, J. Peñuelas^{1,12}, X. Pons², R. Poyatos¹, J. Sardans^{1,12}, O. Sus¹, V. R. Vallejo⁸, J. Vayreda¹ and J. Retana^{1,2}

¹CREAF, Cerdanyola del Vallès 08193, Spain, ²Univ Autònoma de Barcelona, Cerdanyola del Vallès 08193, Spain, ³Univ Granada, iEcoLab Lab Ecol, Instituto Interuniversitario Sistema Tierra, Granada 18006, Spain, ⁴CTFC (Forestry Technology Centre of Catalonia), St Llorenç de Morunys km 2, 25280 Solsona, Spain, ⁵Department of Ecology, University of Granada, E-18071 Granada, Spain, ⁶Department of Biogeography and Global Change, Museo Nacl Ciencias Nat, CSIC (BGC-MNCN-CSIC), E-28006 Madrid, Spain, ⁷Technical School of Agricultural and Forestry Engineering, University of Castilla-La Mancha, Campus Universitario s/n, 02071 Albacete, Spain, ⁸Fundación Centro de Estudios Ambientales del Mediterráneo (CEAM), Paterna, Valencia, Spain, ⁹Department of Biology and Geology, Area Biodiversidad and Conservación, University of Rey Juan Carlos, Mostoles, Madrid, Spain, ¹⁰Department of Biological Sciences, Macquarie University, North Ryde, NSW, Australia, ¹¹CSIC, Institut de Diagnosi Ambiental i Estudis de l'Aigua IDAEA, ES-08034 Barcelona, Spain, ¹²CSIC, Global Ecology Unit CREA-CEAB-UAB, Cerdanyola del Vallès 08193, Spain, ¹³Grupo Interdisciplinar de Sistemas Complejos (GISC), Departamento de Matemáticas, Universidad Carlos III de Madrid, 28911 Leganés, Spain

*Correspondence: Enrique Doblas-Miranda, CREA, Cerdanyola del Vallès 08193, Spain. E-mail: e.doblas@creaf.uab.es

ABSTRACT

Aim Mediterranean terrestrial ecosystems serve as reference laboratories for the investigation of global change because of their transitional climate, the high spatiotemporal variability of their environmental conditions, a rich and unique biodiversity and a wide range of socio-economic conditions. As scientific development and environmental pressures increase, it is increasingly necessary to evaluate recent progress and to challenge research priorities in the face of global change.

Location Mediterranean terrestrial ecosystems.

Methods This article revisits the research priorities proposed in a 1998 assessment.

Results A new set of research priorities is proposed: (1) to establish the role of the landscape mosaic on fire-spread; (2) to further research the combined effect of different drivers on pest expansion; (3) to address the interaction between drivers of global change and recent forest management practices; (4) to obtain more realistic information on the impacts of global change and ecosystem services; (5) to assess forest mortality events associated with climatic extremes; (6) to focus global change research on identifying and managing vulnerable areas; (7) to use the functional traits concept to study resilience after disturbance; (8) to study the relationship between genotypic and phenotypic diversity as a source of forest resilience; (9) to understand the balance between C storage and water resources; (10) to analyse the interplay between landscape-scale processes and biodiversity conservation; (11) to refine models by including interactions between drivers and socio-economic contexts; (12) to understand forest-atmosphere feedbacks; (13) to represent key mechanisms linking plant hydraulics with landscape hydrology.

Main conclusions (1) The interactive nature of different global change drivers remains poorly understood. (2) There is a critical need for the rapid development of regional- and global-scale models that are more tightly connected with large-scale experiments, data networks and management practice. (3) More attention should be directed to drought-related forest decline and the current relevance of historical land use.

Keywords

Assessment, atmosphere-biosphere interaction, biodiversity, climate change, fire regime, forest management, factors interaction, land-use change, research gaps, water reserves.

INTRODUCTION

The earth system is changing, threatening the ecosystem services upon which we depend (Steffen *et al.*, 2004). Greenhouse gas emissions are causing climate change, characterized by warmer temperatures and more frequent and intense droughts (Giorgi & Lionello, 2008), which, in turn, imply an increase in climatic fire risk (Pausas, 2004). Other anthropogenic changes include major changes in land use, increasing nitrogen deposition and the accumulation of tropospheric ozone (Steffen *et al.*, 2004).

Mediterranean terrestrial ecosystems (MTEs), including forests, shrublands and pastures, serve as exemplary natural laboratories in which to study global change because they are highly sensitive to several drivers of such change and to the interactions among these drivers (Sala *et al.*, 2000). The climate in MTEs shows a high sensitivity to global atmospheric changes due to the transitional nature between arid and temperate regions in these ecosystems (Giorgi, 2006). Increased aridity is expected in most existing MTEs (Sillmann *et al.*, 2013). The combination of extreme climate events, a long history of land-use changes and the particular geology of these ecosystems has resulted in more frequent and intense fires, water scarcity and land degradation (soil and productivity loss), among other impacts (Conacher, 1998; Keeley *et al.*, 2012). MTEs show high levels of heterogeneity at different scales due to these disturbances and large seasonal and inter-annual climatic variability (Rundel, 1998). This distinctive spatiotemporal variability of environmental factors has resulted in a singular and diverse biota, with elevated vulnerability to extinction induced by global change (Malcolm *et al.*, 2006). Given that socio-economic trends are projected to have a greater effect than climatic drivers on land use (Schröter *et al.*, 2005), it is important to be able to study global change in different social and economic contexts and across different policy regimes. Among the world's MTEs, the Mediterranean Basin serves as a particularly valuable laboratory for studying global change because of its wide range of socio-economic conditions and government policies (Brauch, 2003).

A consistent evaluation of the principal research gaps within MTEs, with special consideration given to the Mediterranean Basin, has the potential to provide valuable information on how to advance in handling the future impacts of global change on a global scale. A previous study by the International Geosphere-Biosphere Programme (Lavorel *et al.*, 1998; hereafter La98) provided the first roadmap for conducting global change research in Mediterranean Basin ecosystems and recommended that specific actions be carried out in the region. Now, after 15 years of increasing research efforts devoted to the impacts of global change, it is time to evaluate the current state of the art with regard to these targets and to propose a new set of priorities for the coming decades – including priorities relevant to the MTEs outside the Mediterranean Basin.

To do this we revisit the priorities recommended by La98, following their original order, and we provide an update of their current state and relevance. We then suggest a new set of priorities for the future. Our specific objectives are: (1) to evaluate the

progress of the research carried out in MTEs; (2) to assess the accomplishment of previous priorities; and (3) to update the list of priorities with emerging topics and challenges.

EVALUATION OF THE ACCOMPLISHMENT OF THE LA98 RESEARCH PRIORITIES

To understand future fire regimes and their effects

Prediction of future fire regimes, involving interaction with global changes

Land-use change may modify the fire regime by altering fuel load and distribution or ignition patterns. In the MTEs of Europe, massive abandonment of agricultural land has increased landscape homogeneity and crown fire potential, which in turn facilitates the spread of fire (Lloret *et al.*, 2002; Mitsopoulos & Dimitrakopoulos, 2007). In contrast, central Chile has experienced an overall trend of deforestation and loss of shrubland but no clear trend in burned area (Montenegro *et al.*, 2004). In California, urban development is the major driver of land transformation, increasing ignitions in the wildland–urban interface (Syphard *et al.*, 2007).

In the Mediterranean Basin, fire-suppression policies may reduce fire size in the short term but promote megafires in the longer term (Piñol *et al.*, 2007; Brotons *et al.*, 2013). In contrast, fuel reduction policies are regularly implemented with prescribed fires in several MTEs in order to reduce hazard in populated areas and to reproduce a fire regime that supports biodiversity in a fine-grain mosaic of vegetation (Price & Bradstock, 2010). Prescribed burning, however, remains controversial (Boer *et al.*, 2009; Fernandes *et al.*, 2011). Under extreme weather conditions, fuel quantity no longer serves as a control on fire regime and large fires may even occur with relatively low fuel loads (Keeley & Zedler, 2009; San-Miguel-Ayanz *et al.*, 2013). Site idiosyncrasy is also important, affecting pre-fire patch grain and distribution (Syphard *et al.*, 2007). Fire-induced landscape homogenization (by large or frequent fires) as opposed to heterogenization (by scattered fires through space and time) appears highly dependent on fire regime (Lloret *et al.*, 2002).

Climate change is expected to increase dryness in MTEs due to warmer temperatures and reduced precipitation, particularly during the summer (Giorgi & Lionello, 2008; Sillmann *et al.*, 2013). Overall, this tendency will lead to increasing climatic fire risk (Liu *et al.*, 2010), as already apparent from historical records (Kraaij *et al.*, 2013). However, this trend is also influenced by fuel availability, determined by both the quantity of fuel and climate-driven fuel moisture (Westerling *et al.*, 2006; Battlori *et al.*, 2013). Although lower air humidity and fuel water content are positively related to fire ignition and propagation, a drier climate eventually leads to a decrease in fuel load, due to lower productivity (Lenihan *et al.*, 2003; Battlori *et al.*, 2013).

CO₂ fertilization could affect fuel accumulation and thus fire regimes. Recent projections at the regional level reveal that the net effect of CO₂ fertilization combined with drought stress

remains uncertain (Keenan *et al.*, 2011). However, water appears to be more important than CO₂ as a driver of growth in water-limited MTEs (Faticchi *et al.*, 2014). In addition, the combination of increased fire frequency and drought stress could enhance shrub encroachment (Pausas, 1999; Mouillot *et al.*, 2002) and thus reduce carbon storage.

Fire impacts

On landscape patterns

The main aim behind the extensive development of spatially explicit landscape models for MTEs is to assess the vulnerability of forests to fire due to soil degradation or vegetation shifts (Franklin *et al.*, 2005; Millington *et al.*, 2009). Most of these models, therefore, include forest management and environmental factors within specific landscape configurations (Loepfe *et al.*, 2011; Moreira *et al.*, 2012). However, models are still incomplete, in the sense that anthropogenic factors are usually not included in the generation of landscape projections and thus the prediction of future fire impacts (Syphard *et al.*, 2007; Le Page *et al.*, 2010).

On vegetation

Simulation models of fire and vegetation dynamics, including the interaction between these variables, have been developed, considering different spatial scales, vegetation levels, successional approaches and explicit or implicit simulation of fire spread (Keane *et al.*, 2004; Millington *et al.*, 2009). Analysis of the regenerative traits of species and populations remains a key approach in the assessment of the sensitivity of plant species to predicted fire regimes and for modelling changes in species distribution and community composition (Lloret *et al.*, 2005; Syphard & Franklin, 2010). Several studies have assessed the short-term responses of MTE vegetation (Keeley *et al.*, 2012; Moreira *et al.*, 2012) but more attention should be paid to longer time frames.

Combined with other factors

The consequences of the combined effects of fire and drought in the form of a decline in the regeneration of plants and land degradation have been the object of intense study (e.g. Mouillot *et al.*, 2002; Montenegro *et al.*, 2004). More recently, the role of fire in facilitating the spread of biological invasions in MTEs has also attracted attention (Rouget *et al.*, 2001; Pino *et al.*, 2013) and research in this area is revealing the vulnerability associated with properties of the fire regime such as fire frequency (Keeley & Brennan, 2012) and intensity (Franklin, 2010).

Fire control and mitigation

Prevention

Since La98, many studies have been conducted to characterize the relationship between different fire spread characteristics using diverse approaches. These include fire behaviour models

coupling suppression policies and vegetation dynamics, forest inventories, wildfire databases and fire severity studies (e.g. Boer *et al.*, 2009; Price & Bradstock, 2010). Such studies have provided quantitative information about the best way to manage forest fuels to obtain: (1) more favourable fire characteristics (behaviour, frequency, size) aiding future fire suppression (Crecente-Campo *et al.*, 2009) or unplanned fire extents (Boer *et al.*, 2009), and (2) landscape configurations that are more efficient at reducing megafires (Millington *et al.*, 2009; Loepfe *et al.*, 2011). Several studies have also reported positive feedback between flammable vegetation types and fire frequency (Vilà *et al.*, 2001; Grigulis *et al.*, 2005).

Restoration

The last decade has seen substantial advances in key aspects of post-fire forest restoration, such as the analysis of vegetation recovery (Díaz-Delgado & Pons, 2001; Díaz-Delgado *et al.*, 2002). A new field of study deals with the identification of previous land-use changes as drivers of post-fire regeneration (Clavero *et al.*, 2011; Puerta-Piñero *et al.*, 2012). Additional breakthroughs have been made in the study of facilitative interactions and the use of shrubs as potential nurse plants (Gómez-Aparicio *et al.*, 2004), underscoring the role of early and mid-successional shrubs for forest regeneration in burnt areas (Gómez-Aparicio *et al.*, 2004; Siles *et al.*, 2010). In addition, a novel focus has recently appeared in relation to the use of coarse woody debris to foster restoration success, as this debris may act as a nurse structure, improving microclimatic conditions for seedling establishment, increasing soil nutrient content and improving other physical and chemical properties of the soil (Castro *et al.*, 2011; Marañón-Jiménez & Castro, 2013; Marzano *et al.*, 2013). Burnt trees are also a biological legacy crucial for the recovery of communities and for the structure and function of regenerating mediterranean-type ecosystems by promoting plant-animal interactions and increasing plant and animal diversity (Castro *et al.*, 2012; Lee *et al.*, 2013; Marzano *et al.*, 2013; Leverkus *et al.*, 2014), reducing invasion by exotic species (Moreira *et al.*, 2013), promoting soil microbial activity (Marañón-Jiménez & Castro, 2013) and carbon sequestration (Serrano-Ortiz *et al.*, 2011).

To study the effects of land use on biosphere–atmosphere interactions

Feedbacks of land-use changes on the climate system

While a great deal of uncertainty persists concerning the impacts of land-use changes on regional climate models (Bonan, 2008), modelling results for the Mediterranean suggest that changes in land use significantly affect climate (Lionello *et al.*, 2006). Changes in evapotranspiration rates and surface albedo due to deforestation in the Mediterranean Basin could provoke cooler and moister springs but warmer and drier summers (Heck *et al.*, 2001) or instead cooling during summer (Zampieri & Lionello, 2011). In south-west Australia, deforestation may lead to long-term reductions in rainfall patterns (Pitman *et al.*, 2004). In

summary, available models indicate that the climate of MTEs is sensitive to changes in vegetation cover, especially in summer. However, the induced climate anomalies can be associated with fine-grained, complex and non-local mechanisms and their relative weights in driving local effects are still uncertain (Seneviratne *et al.*, 2010).

Feedbacks of ecosystem physiology on the climate system

Current evidence suggests that CO₂-driven feedbacks on temperature through physiological responses of vegetation are negligible (Keenan *et al.*, 2011; Cheaib *et al.*, 2012).

The effects of climate change on soil respiration and the emission of other biogenic gases are still a major concern. Increases in drought intensity in the Mediterranean Basin have been associated with episodes of tree mortality during recent decades (Martínez-Vilalta *et al.*, 2012; Sánchez-Salguero *et al.*, 2012). A recent study shows that although current tree mortality is not affecting the carbon balance of Mediterranean forests, increased warming and drought are likely to alter the capacity of these forests to absorb CO₂ and forest management may be a key factor determining the response of forest C balance to changing climate (Vayreda *et al.*, 2012).

Some studies have reported an increase in soil organic carbon (SOC) and other nutrient fractions under drought due to rising quantities of litterfall and dead roots (Talmon *et al.*, 2011) and to decreased soil decomposition and respiration (e.g., Curiel Yuste *et al.*, 2007; Ryals & Silver, 2013). However, observational studies suggest that drought should decrease SOC in the long term by reducing plant cover, which implies a decrease in litterfall, soil protection and permeability (Boix-Fayos *et al.*, 1998). Furthermore, in MTEs there is an overall reduction in biological activity in the soil due to a reduction in soil moisture (Brown *et al.*, 1996), and this results in a decrease in soil nutrient availability and soil CO₂ emissions (Emmett *et al.*, 2004; Sardans *et al.*, 2008). An increasing body of research is also starting to reveal the key role of microbial communities in ecosystem processes under climate-change scenarios, particularly in soil carbon dynamics (Balsler & Wixon, 2009; Asensio *et al.*, 2012; Curiel Yuste *et al.*, 2012).

Biogenic volatile organic compounds (BVOCs) were not previously considered but are also crucial for understanding both biological consequences and feedbacks on atmospheric chemistry and climate itself (Monson *et al.*, 2007; Peñuelas *et al.*, 2013). The emission rates of BVOCs increase with rising global temperature, but changes in species and community structure, land use and resource availability can also lead to major changes in Mediterranean regional BVOC fluxes (Peñuelas & Staudt, 2010).

Contribution of fire related emissions of carbon, nitrous oxides and other trace gases to the atmosphere and their potential effects on climate

Fire emissions cause significant perturbations of the chemical composition of the atmosphere and in the earth climate system, as several field campaigns as well as laboratory experiments and

prescribed burnings in MTEs have shown (Ciccioli *et al.*, 2001; Phuleria *et al.*, 2005; Wain *et al.*, 2008; Garcia-Hurtado *et al.*, 2013). The principal gases emitted include carbon dioxide (CO₂), carbon monoxide (CO) and methane (CH₄), but also nitrogen oxides (NO_x and N₂O), ammonia (NH₃), sulphur dioxide (SO₂), light hydrocarbons, volatile and semi-volatile organic compounds, and particulate matter (10–2.5 µm), which could affect climate change and human health (Ciccioli *et al.*, 2001; Bell & Adams, 2008).

Studies have been conducted to characterize biomarkers from woodstove combustions and wildfires (Muhle *et al.*, 2007; Gonçalves *et al.*, 2011), but the challenge of identifying the main tracer compounds emitted by forest fires in MTEs continues.

EMEP/CORINAIR emission inventories and satellite observations, including those using Moderate Resolution Imaging Spectroradiometer (MODIS) data, have been used to feed several emission and air quality models (Lazaridis *et al.*, 2005; Paton-Walsh *et al.*, 2012).

Coupled biosphere–atmosphere models for the Mediterranean Basin

Large-scale generalized simulations of the effects of climate and land-use changes on MTEs exist (Zaehle *et al.*, 2007), but regional applications to key ecosystems (including cropland phenology and management to account for associated albedo feedbacks; Sus *et al.*, 2010) are scarce. During the past decade, the development of non-fully-coupled models simulating Mediterranean terrestrial biosphere–atmosphere interactions has focused on the integration of information on ecosystem physiology and land-cover dynamics (e.g. Gritti *et al.*, 2006). Only in recent years, however, have fire models begun to be incorporated into land-surface models (e.g. Prentice *et al.*, 2011). Early efforts identified model deficiencies in reproducing the response of leaf gas exchange to drought events (Reichstein *et al.*, 2003), leading to subsequent model development (Garbulsky *et al.*, 2008; Keenan *et al.*, 2010). Despite this, models continue to perform poorly in conditions of water stress (Vargas *et al.*, 2013). Many ecosystem disturbances, processes and physiological responses remain poorly understood, and are not explicitly accounted for in models, such as, for example, competitive interactions, the depletion of carbohydrate reserves and stress-induced plant decline and mortality (e.g. Carnicer *et al.*, 2011; Misson *et al.*, 2011).

To study landscape effects on water availability and quality

Research at the patch scale

Leaf area index (LAI) and hydrological response

Land-use changes, hydroclimatic conditions and fires are the main drivers of changes in vegetation cover in MTEs and therefore exert great influence on the hydrological responses of these ecosystems. Fires, for instance, influence understorey regrowth

and hence may contribute to maintaining ecosystem evapotranspiration fluxes (Macfarlane *et al.*, 2010). Land-use history also modulates the hydrological behaviour through changes in soil properties such as soil water repellency and infiltrability (Llovet *et al.*, 2009).

Drought-induced episodes of vegetation dieback may result in different ecohydrological effects compared with canopy cover changes (Adams *et al.*, 2012). For example, generalized drought-induced defoliation across southern European forests (Carnicer *et al.*, 2011) may gradually reduce stand transpiration (e.g. Limousin *et al.*, 2009). Drought severity and/or duration together with local factors such as soil water-holding capacity (Peterman *et al.*, 2013) and species-specific drought-tolerance traits (Jacobsen *et al.*, 2007; Matias *et al.*, 2012) will ultimately determine plant survival and, hence, the impact of episodic drought on hydrological processes.

Temporal variability

In general, evapotranspiration in MTEs is strongly depressed during summer (Baldocchi *et al.*, 2010; Raz-Yaseef *et al.*, 2012). However, mediterranean plant species in MTEs show a variety of responses to cope with drought, from complete summer senescence observed in some grasslands (Baldocchi *et al.*, 2004) to various degrees of stomatal control of transpiration (e.g. Quero *et al.*, 2011). In general, increased atmospheric CO₂ concentrations will induce stomatal closure and improve plant water status, enhancing water-use efficiency and potentially reducing the effects of increases in evaporative demand. However, the impact of increased CO₂ on ecosystem water use will vary with vegetation type, species and stand development (Li *et al.*, 2003). Regardless of all these functional responses to water deficits, water (and carbon) fluxes in MTEs are strongly reduced during extreme drought events (Granier *et al.*, 2007).

Hydrological equilibrium and simulation models

In MTEs, maximum LAI is constrained by vegetation type, local climate and soil conditions, leading to the notion that there is an 'equilibrium LAI' that maximizes carbon assimilation (Hoff & Rambal, 2003). This hypothesis was framed by La98 within Eagleson's (1982) broader concept of hydrological equilibrium, but the optimality hypotheses associated with Eagleson's ecohydrological model have now been questioned in the context of water-limited environments (Kerkhoff *et al.*, 2004).

Research at the landscape scale

Mapping and emergent properties

The availability of earth observational data for ecological studies has increased due to the launch of several multispectral high/medium spatial resolution platforms and it will increase further in the coming years with Landsat-8 and ESA's Sentinel missions. These developments will give continuity to the wide use of remote sensing in previous MTE studies (Shoshany, 2000). Lidar

technology is becoming an operative application for acquiring information about forest and shrubland structures in many regions (Estornell *et al.*, 2010; García *et al.*, 2010). In contrast, radar information, a promising data source in the past, has not provided the expected results (Lu, 2006). MTEs have been increasingly studied using remote sensing from different perspectives, including land-use/cover changes, drought, carbon budget and foliar biochemical concentration and canopy structure (e.g. Serrano *et al.*, 2001; Berberoglu & Akin, 2009). Unmanned aerial systems (UAV) provide another increasingly popular platform for ecological studies (Dunford *et al.*, 2009; Hernández-Clemente *et al.*, 2012). Another area of interest is the use of spectroradiometers as an augmentation to remote sensing for validating or training models (Xu & Baldocchi, 2004; Glenn *et al.*, 2011).

Landscape change

Different studies at the operational catchment scale in several MTEs have evidenced decreasing trends in the flow records (e.g. Delgado *et al.*, 2010; Zhao *et al.*, 2010; see also Lespinas *et al.*, 2010, for contrasting results) and modifications of the flow regime (e.g. López-Moreno *et al.*, 2011; Morán-Tejeda *et al.*, 2011), partly (but not exclusively) attributable to forest expansion. The effects of climate change on surface water quality and ecology are attracting growing attention in mediterranean areas (e.g. Munné & Prat, 2011; Otero *et al.*, 2011). Investigations on post-fire flows of soil, water and nutrients, however, have not been so common, mainly because pre-fire data are frequently unavailable (Shakesby, 2011) and the effects of prescribed fires do not directly mimic natural wildfire influences (Seibert *et al.*, 2010).

Simulation models

Numerous complex hydrological models (often spatially distributed, physically based) have been used to assess the effects of global change on water resources, soil erosion and vegetation productivity (e.g. D'Agostino *et al.*, 2010; Senatore *et al.*, 2011). However, there is growing concern about the current modelling approaches (Ewen *et al.*, 2006; Beven, 2011) because of cumulative uncertainties in regional model projections, including uncertainties in downscaling procedures, land-use and land-cover changes and choices of hydrological models.

To investigate the effects of climate change on ecological diversity

Genetic diversity

La98 suggested giving increased attention to studies linking genetic diversity with relevant ecosystem functions. However, there is still little knowledge of the adaptive variability present in the different MTEs. The technical difficulty of some measurements, the time-consuming nature of the relevant experiments and the large sample sizes required to determine heritability or

selection on functional traits have limited the implementation of reciprocal transplants and/or common garden studies, particularly for long-lived species (Ackerly, 2006). Exceptions include economically important species for which provenance trials have allowed the estimation of ecotypic variation in growth and functional traits (e.g. Ramírez-Valiente *et al.*, 2009; Kurt *et al.*, 2012).

Significant advances have been made in the last decade in understanding the range dynamics of mediterranean species in response to past climatic changes to validate predictions on ecological and evolutionary consequences of current climate change (e.g. Petit *et al.*, 2005, 2008). In addition, there has been mounting evidence showing that maintaining genetic diversity within natural populations can maximize their potential to withstand and adapt to biotic and abiotic disturbances (Jump *et al.*, 2008).

Species and functional diversity

Since La98 noted the effects of diversity on ecosystem functioning (DEF), those effects have continued to be intensely discussed in MTEs, although the relative importance of DEF in relation to global change still needs to be further assessed (Larsen *et al.*, 2005; Dimitrakopoulos, 2010; Maestre *et al.*, 2012). Further, the role of biodiversity in maintaining human-made systems that provide high-value ecosystem services is becoming a key issue in shaping environmental and land-use policies (Díaz *et al.*, 2013).

In the Mediterranean Basin, higher diversity has been associated with more rapid recovery after fire (Lavorel, 1999). Fire effects also interact with species richness, increasing biomass production in rich communities (Dimitrakopoulos *et al.*, 2006). In most MTEs, restoration of degraded environments due to human alteration could be facilitated by diversity of vegetation and associated soil microbes (García-Palacios *et al.*, 2011; Viers *et al.*, 2012). Recent work also emphasizes the role of diversity in the face of invasive species (Prieur-Richard & Lavorel, 2000; Selmants *et al.*, 2012; see also Prieur-Richard *et al.*, 2002, for contrasting results). Also, the effect on ecosystem functioning of invasion-induced species impoverishment is a contested issue (Vilà *et al.*, 2006; Ruwanza *et al.*, 2013).

Aboveground–belowground trophic interactions are of potential importance in the functioning of MTEs (e.g. Doblas-Miranda *et al.*, 2009; Janion *et al.*, 2011). Several experiments in mediterranean regions have related soil respiration and functioning to climate and land-use change (Emmett *et al.*, 2004; Garnier *et al.*, 2007; Lau & Lennon, 2012).

The expected increase in aridity in most MTEs may impact plant community dynamics and composition (Lloret *et al.*, 2009; Matías *et al.*, 2012). However, the heterogeneity of mediterranean landscapes and the variety of responses at different scales give rise to a variety of stabilizing mechanisms promoting community resilience (Lloret *et al.*, 2012) and make predictions difficult (Maestre *et al.*, 2005). Climate change could also affect the biodiversity of important and influential faunal communities (Botes *et al.*, 2006; Gil-Tena *et al.*, 2009). Another promising line of research is to disentangle the combined effects of

different factors of change on species and functional diversity (Gil-Tena *et al.*, 2009; Pasquini & Vourlitis, 2010).

Landscape diversity

The increasing availability of geospatial information at increasing spatial and temporal resolution has facilitated the study of landscape patterns. Projects in the Mediterranean Basin, in California's mediterranean landscapes and in South Africa have boosted the development of studies addressing how drivers of global change might influence diversity and key ecosystem functions at broad spatial scales (e.g. Santos *et al.*, 2006; Aparicio *et al.*, 2008; García *et al.*, 2011).

Current research is demonstrating that landscape structure modulates conservation efforts at local scales due to non-linear effects of landscape diversity on local diversity (Concepción *et al.*, 2012). Conservation at local scales would be effective only at intermediate levels of landscape complexity, whereas landscape initiatives would be more effective in the simpler and more complex landscapes (Brotons *et al.*, 2004; Concepción *et al.*, 2008). Landscape-scale management is crucial for preserving both biodiversity and the ecosystem services the landscape provides (Díaz *et al.*, 2013).

The effects of landscape heterogeneity also have a strong temporal dimension. Due to the long history of land-use changes in the Mediterranean Basin, there is increasing evidence of long-term impacts of past land uses on the current state of ecosystems (Puerta-Piñero *et al.*, 2012; Navarro-González *et al.*, 2013). In this way, several projects have aimed to disentangle the role of past land uses at different scales and in different ecosystems (Bonet *et al.*, 2010; Ortega *et al.*, 2010).

UPDATE OF PRIORITIES AND NEW CHALLENGES

Advancement of research effort in MTEs

In general, the study of global change in mediterranean ecosystems has increased since 1998. The proportion of global change studies on MTEs remains relatively low, but is now closer to the proportion of global change studies devoted to other ecosystems, which have diminished (boreal, tropical) or remained approximately constant (temperate) (Fig. 1). There has been an increase in studies in all four lines of research proposed by La98, especially from the mid 2000s, and particularly in the last few years (Fig. 2). This increase is quite similar for fire regimes, water availability and quality and ecological diversity, and somewhat lower for biosphere–atmosphere interactions. Studies about the effects of global change in mediterranean ecosystems have been mostly carried out in the Mediterranean Basin (88.4% of total studies), while studies within this region outside European countries (i.e. the southern rim of the Mediterranean Basin) constitute just a small fraction (7.8% of the Mediterranean Basin studies).

Figure 1 Trends in published research articles (excluding reviews and meeting abstracts) related to global change and mediterranean forests in the scientific literature, for the years 1998–2012, based on ISI Web of Science search. The term forest, but not shrublands or pastures, has been used for comparative purposes. Plotted lines show the percentage of references retrieved using the topic words ‘(forest change) AND (fire or atmospher* or ‘land use’ or water or diversity) NOT (marine sea)’ plus a regional definition (mediterranean, boreal, tropical or temperate), relative to all references obtained without the regional specification.





Figure 2 Trends in published research articles (excluding reviews and meeting abstracts) related to global change and forests in the scientific literature, for the years 1998–2012, based on ISI Web of Science search. The term forest, but not shrublands or pastures, has been used for comparative purposes. Plotted lines show the total of references, after a one by one selective review to avoid articles not really related with the subject, using the following topic words.

For fire regimes: (mediterranean forest fire change) AND (predict* OR model OR simulation OR impact OR effect OR consequence OR control OR prevention OR restoration) AND ('land use' OR 'land cover' OR landscape OR climate OR temperature OR humidity OR moisture OR atmospher* OR plant OR vegetation OR 'ecosystem processes') NOT (marine sea).

For biosphere–atmosphere interactions: (mediterranean forest atmospher* change) AND (feedback OR model OR soil OR physiology OR biogenic OR volatile OR emissions OR monitoring OR carbon OR 'trace gases' OR 'nitro* oxides' OR ammonia OR sulphur OR ozone OR transpiration OR eddy-covariance) AND ('land use' OR 'land cover' OR landscape OR climate OR temperature OR humidity OR water OR energy OR moisture OR fire OR plant OR tree OR vegetation OR 'ecosystem processes') NOT (marine sea).

For water availability and quality: (mediterranean forest water change) AND (availability OR quality OR model OR simulation OR flow OR flux OR 'patch scale' OR 'landscape scale' OR 'leaf area' OR hydrological OR 'temporal variability' OR trend OR mapping OR 'remote sensing' OR catchment) AND ('land use' OR 'land cover' OR landscape OR climate OR temperature OR humidity OR moisture OR atmospher* OR fire OR plant OR vegetation OR 'ecosystem processes') NOT (marine sea).

For ecological diversity: (mediterranean forest diversity change) AND (bioindicator OR conservation OR restoration OR heterogeneity OR trait OR genotyp* OR phenotyp* OR model OR simulation OR impact OR effect OR trophic OR genetic OR species OR functional OR ecological OR biological) AND ('land use' OR 'land cover' OR landscape OR climate OR temperature OR humidity OR moisture OR fire OR invasi* OR atmospher* OR soil OR 'ecosystem processes' OR 'ecosystem function*') NOT (marine sea).

Table 1 Overview of the degree of accomplishment of the priorities of La98 by the scientific community. It should be noted that while practical for abstracting purposes a categorical ‘yes’ or ‘no’ is never applied to science. We recommend therefore that ‘yes’ is considered as a ‘great effort invested in this particular subject’ and ‘no’ as ‘although some attempts have been made to study the subject, it still needs further development’.

Previous priorities		Accomplishment quick view		
1. Understand future fire regimes	1.1 Prediction	Effects of land use	Partially	
		Effects of climate	Yes	
		Effects of atmospheric composition	No	
	1.2 Impacts	On landscape	Yes	
		On vegetation	Partially	
		Combined with climate change	Yes	
		On ecosystem processes	Partially	
	1.3 Control	Prevention	Yes	
		Restoration	Partially	
2. Study biosphere–atmosphere interactions	2.1 Land use and climate	Partially		
	2.2 Physiology and climate	CO ₂ and temperature	Partially	
		Temperature and biogenic emissions	Partially	
	2.3 Fire emissions	Yes		
3. Landscape effects on water	2.4 Coupled models		Partially	
		3.1 Patch scale	LAI and hydrological response	Yes
			Hydrological equilibrium	Partially
			Temporal variability	Partially
		Simulation models	No	
	3.2 Landscape scale	Mapping	Yes	
		Scales and emergent properties	Yes	
Landscape change		Yes		
4. Changes in ecological diversity		Simulation models	Partially	
	4.1 Genetic	Partially		
	4.2 Species and functional	Partially		
	4.3 Landscape	Partially		

LAI, leaf area index.

New research priorities

Since 1998, great advances have been made in research techniques and knowledge. Most of the research priorities identified by La98 have been addressed (Table 1), but some remain and new MTE research topics and priorities have also emerged. Our evaluation has identified: (1) topics listed in La98 which have been only partially addressed, (2) new approaches to questions already suggested by La98, and (3) a new set of emerging issues (Table 2). We suggest a new classification of these priorities based on a framework describing how Mediterranean ecosystems will respond to human-induced global change, including ecosystem services derived from ecosystem functioning, and incorporating mechanisms of monitoring, studying and seeking to modify these responses (Fig. 3).

In our proposed scheme, the first step in organizing future research efforts on global change in MTEs (and other ecosystem types) is to understand the effect of the drivers of global change on ecosystem functioning. Second, these processes must be monitored, and data must be appropriately analysed to produce useful research outputs. Third, this information should be used to guide ecosystem management aimed at modifying the observed or expected effects. Fourth, the link between ecosystem

functioning and ecosystem services should be made explicit, in order to fully realize the opportunities offered by ecosystem management in a global change context. Finally, all the previous steps will depend on spatial and temporal scales of functioning, observation and management, which should be analysed by means of ecosystem modelling approaches and well-designed critical experiments.

Effects of the interactions between the drivers of global change on ecosystem functioning

To establish the role of the landscape mosaic on fire-spread

A major challenge remains to establish how direct and indirect fire suppression policies influence fire regimes. These policies operate in spatially explicit landscapes that are determined by fuel load and continuity, which in turn determine fire regime. Specifically, we need to better understand the fraction and distribution of agricultural land needed to prevent the spread of megafires, and the role of the critical wildland–urban interface, as well as the associated modification of landscape structure, in preventing the massive crown fires that can cause megafires (Keeley *et al.*, 2012; Loepfe *et al.*, 2012). In addition, the disrupt-

Table 2 Proposed research priorities for mediterranean terrestrial ecosystems in the face of global change.

New priorities

Effects of the interactions between drivers of global change on ecosystem functioning

- 1 To establish the role of the landscape mosaic on fire spread
- 2 To further research the combined effect of different drivers on biological invasions and pest expansion
- 3 To address the interaction between drivers of global change and recent forest management practices

Monitoring and data assessment of ecosystem response to global change

- 4 To obtain more realistic information, at larger temporal and spatial scales, of the impacts of global change and ecosystem services to be used in models

- 5 To assess forest mortality events associated with climatic extremes (particularly drought)

Managing ecosystems to enhance resilience

- 6 To focus global change research on identifying and managing vulnerable areas
- 7 To use the functional and life-history traits concepts to study resilience and community assembly after disturbance
- 8 To promote cross-disciplinary research to study the relationship between genotypic and phenotypic diversity as a source of forest resilience

Embracing the link between ecosystem functions and services

- 9 To understand how forest management affects the balance between C storage and water resources at large spatial and temporal scales
- 10 To analyse the interplay between landscape-scale processes and biodiversity conservation along wide gradients of landscape complexity

Scaling ecosystem dynamics in space and time under different scenarios

- 11 To refine predictive models by including interactions between global change drivers and socio-economic contexts
- 12 To use manipulative, interdisciplinary and multiscale experiments to understand forest–atmosphere feedbacks
- 13 To improve the representation of key mechanisms linking plant hydraulics with landscape hydrology




Figure 3 Framework for global change research priorities in mediterranean terrestrial ecosystems. The framework is structured according to different causal and observational pathways, as follows: (1) effects of the interactions between global change drivers on ecosystem functioning; (2) monitoring and data assessment of ecosystem response to global change; (3) managing ecosystems to enhance resilience; (4) embracing the link between ecosystem functions and services; (5) scaling ecosystem dynamics in space and time under different scenarios.

tion of the landscape–fire interaction by extreme climatic episodes should be included in these analyses, and increasing vulnerability to fires should be addressed in areas where climate is becoming similar to that of the existing Mediterranean regions.

To further research the combined effect of different drivers on biological invasions and pest expansion

Since La98 did not address the impact of fire in combination with factors other than climate change and land use, it did not include biological invasions. The effect of biological invasions in combination with different climatic events (e.g. droughts), disturbance (e.g. fires) and landscape structure (e.g. the wildland–urban interface) on biodiversity and ecosystem functioning stands as a priority for future research efforts. For example, the impact of the combination of fire and invading biota may be especially important in mediterranean regions where the spread of invasive species is altering fire regimes. In addition, it is important to assess the influence of climate change combined with land-use change in the expansion of certain pest species (native or not) in previously non-accessible habitats like mountains. This new focus will complement the more classical, but still much needed, analyses on how responses of keystone species or communities to global change may ameliorate or amplify direct effects on forest structure and function (e.g. Valladares *et al.*, 2013).

To address the interaction between drivers of global change and recent forest management practices

Human influences have shaped the current structure and composition of MTEs and their woodlands. In some regions these impacts have built up over the last decades as a result of changes in forest management and land-use practices. In the Mediterranean Basin, for instance, there is a widespread process of forest

densification owing to widespread abandonment of intensive forest management. This process has critical implications in a global change context, as it affects fire spread and recovery after fire (e.g. Puerta-Piñero *et al.*, 2012) and increases the competition for water and therefore the likelihood of drought-induced forest die-off (Martínez-Vilalta *et al.*, 2012). Although the demographic implications of the interaction between changes in climate and competition are starting to be addressed (Vilà-Cabrera *et al.*, 2011; Ruiz-Benito *et al.*, 2013) there is an urgent need to expand these studies in order to disentangle the contribution of different drivers (and their interaction) to current stand dynamics and to translate this information into credible models of future forest dynamics.

Monitoring and data assessment of ecosystem response to global change

To obtain more realistic information, at larger temporal and spatial scales, of global change impacts and ecosystem services to be used in models

La98 advised that, in order to disentangle the effects of landscape change on hydrological properties, research on larger spatial scales is necessary. In fact, other ecosystem services such as carbon storage are also better defined and studied at larger spatial scales and over longer time-scales (e.g. Vayreda *et al.*, 2012). Similar reasoning can be applied to the factors that alter these services. For example, there is a need for reliable information on the efficiency of different fire and fuel management alternatives at reasonably large spatiotemporal scales. Long-term manipulative experiments show that ecosystem responses to disturbance frequently change over time (e.g. Barbeta *et al.*, 2013). Thus, the data inputs for model calibration and validation should include, to the extent possible, long time series of observational data as well as long-term ecosystem manipulation experiments (Ecotron) focusing on key drivers (cf. Beier *et al.*, 2012).

To assess forest mortality events associated with climatic extremes (particularly drought)

The drought-induced forest decline detected during recent decades remains insufficiently understood, as it is not yet known which biological mechanisms are involved and which factors other than drought have played causal roles. In MTEs, elevation, substrate, plant composition, stand structure and soil biota all appear to contribute to forest die-off (e.g. Martínez-Vilalta *et al.*, 2012). Forest history and, particularly, management, appear to be key drivers, for instance by determining current forest structure and composition. Strategic actions include long-term monitoring at a regional scale, with implementation of common protocols, and rapid identification of new events (using for example remote sensing or UAVs), which would make it possible to study the process while it is happening. These actions would benefit critically from the involvement of forest owners and governmental agencies.

Managing ecosystems to enhance resilience

To focus global change research on identifying and managing vulnerable areas

Future research efforts should aim at identifying areas that might suffer from the combination of multiple drivers of climate change. For instance, mountains and sub-mediterranean zones may be particularly susceptible to the likely increase of climatic fire risk because these are landscapes that have not previously faced high fire risks and therefore may have low vegetation resilience (Lloret *et al.*, 2005). Other susceptible areas include those where land-use transformation has led to a high fuel load and thus a high risk of massive crown fires. Similarly, the recent increase in temperature is likely to favour insect expansion, and unprecedented insect outbreaks can arise in areas with high tree density and landscape connectivity. The resulting management agenda should include the adaptation of MTEs to more arid conditions, for instance by species selection, or by management of stand and landscape structure and water use. Here, one major challenge is to determine how the suppression of wildfires and other management actions could eventually induce non-reversible state transitions of vegetation types and structure, resulting in service-impoverted states.

To use functional and life-history traits concepts to study resilience and community assembly after disturbance

Although the possibility of threshold-type responses in ecosystems is real and should be taken into account, the resilience of ecosystems to climate change is frequently substantial and deserves further study (cf. Lloret *et al.*, 2012). Given the variability of species responses to different disturbance types (e.g. the responses of plants to fire regimes; Pérez *et al.*, 2003; Rey Benayas *et al.*, 2007) and the complexity of the factors shaping community dynamics there is an urgent need to find synthetic, yet powerful approaches to predict the ecosystem-level effects of environmental changes. In that respect, the functional trait concept has strong conceptual appeal (Lavorel & Garnier, 2002), although its empirical applicability remains to be properly established.

To promote cross-disciplinary research to study the relationship between genotypic and phenotypic diversity as a source of forest resilience

Given the complexities underlying evolutionary processes, there is a clear need to intensify cross-disciplinary research among different disciplines such as genetics, genomics, demography, functional ecophysiology and animal-plant interactions in order to investigate the effects of climate change on genetic diversity. Reciprocal transplants and common garden experiments, linked to next-generation sequencing approaches, have yet to be fully applied in mediterranean contexts. To understand how mediterranean species will respond to global change in the long term and what the genetic basis of this response will be, it

is essential to identify genes under natural selection, as well as to ascertain the relationship between naturally occurring genotypic and phenotypic diversity.

Embracing the link between ecosystem functions and services

To understand how forest management affects the balance between C storage and water resources at large spatial and temporal scales?

It has been recently suggested that management could increase the C storage capacity of forests (Vayreda *et al.*, 2012), although the detailed mechanisms are yet to be properly characterized. Indeed, when assessing the C absorption capacity of MTE forests it is necessary to consider the importance of the water balance; tree density and species should be managed cautiously since high transpiration rates also imply high water losses in the system. The search for management practices favouring C storage should take into account the risk of forest decline due to scarcity of water.

To analyse the interplay between landscape-scale processes and biodiversity conservation along wide gradients of landscape complexity

Overall, large-scale collaborative efforts among teams in different regions should be promoted and prioritized in order to fully understand how and why landscape processes influence the responses of MTEs to global change. Biodiversity is likely to be crucial in modulating such responses, so effective conservation strategies over wide gradients of landscape complexity need to be designed and effectively implemented. In addition, research on the social and economic role of biodiversity in the maintenance of land-use systems is urgently needed (Campos *et al.*, 2013) in order to establish how and why biodiversity is contributing to the ecological and economic sustainability of mediterranean low-intensity management systems of high natural value.

Scaling ecosystem dynamics in space and time under different scenarios

To refine predictive models by including interactions between drivers of global change and socio-economic contexts

More attention should be given to anthropogenic factors when generating landscape projections (Serra *et al.*, 2008), to understand not only future fire impacts (Brotons *et al.*, 2013) but also other components of global change such as biological invasions and land-use changes, taking into account interactions between these major drivers and the impacts on associated ecosystem services (Campos *et al.*, 2013). Among these factors, further research needs to focus on the integration of the socio-economic scenarios that are already available into predictive models of land-use-driven climatic change (Verburg *et al.*, 2010). The development of reference scenarios of land-use and forest change

that can be used consistently together with available climate change scenarios remains a gap in current global change science. Furthermore, even if the present review is focused on forests, shrublands and pastures, the simulation of biosphere-atmosphere interactions in other key ecosystems such as croplands and urban environments should be considered.

To use manipulative, interdisciplinary and multiscale experiments to understand forest-atmosphere feedbacks

La98 were already conscious that manipulative experiments at large scales are needed to fully understand the feedbacks between terrestrial ecosystems and the atmosphere, in addition to a wider use of historical data. We have learned from a first generation of manipulative field experiments and we are now in a position to use new designs that are larger in scope and make full use of research networks working with standardized protocols (see the review by Beier *et al.*, 2012, for precipitation manipulation experiments). Multifactorial (e.g. CO₂ × warming × drought) and interdisciplinary experiments combining different experimental approaches (field and microcosm experiments) with the use of innovative techniques (e.g. genome pyrosequencing, solid-state nuclear magnetic resonance) will facilitate the identification of key mechanisms and their integration into predictive models. Emerging issues, such as altered BVOC emissions, nutrient imbalances and soil microbial processes, deserve special attention.

To improve the representation of key mechanisms linking plant hydraulics with landscape hydrology

There is a need for improved representations of soil and plant hydraulics in process models of water and carbon fluxes, both in general and for mediterranean vegetation in particular (e.g. Hernández-Santana *et al.*, 2009). Mechanisms leading to drought-induced vegetation die-off are poorly understood and their representation in models is frequently inadequate (McDowell *et al.*, 2013), which limits our capacity to predict vegetation shifts and corresponding ecosystem-level implications (Anderegg *et al.*, 2013). In addition, hydrological predictions are complicated by the difficulty in properly accounting for 'natural' successional dynamics and predicting stochastic events like insect outbreaks and the occurrence of fire.

BROAD RECOMMENDATIONS

From the full list of priorities offered in Table 2, three broad recommendations serve as a conclusion.

1. The interactive nature of different global change drivers remains poorly understood. Different global change factors and their interactions, as well as socio-economic constraints, must be included in the forecasts and modelling of future ecosystem changes.
2. There is a critical need for rapidly developing regional- and global-scale models. Better networking of research data, as well

as manipulative experiments, covering different abiotic and biotic factors and different temporal and spatial scales, is needed to calibrate and validate these ecological models.

3. More attention should be directed at emerging issues especially related to MTEs in the face of global change, including recent drought-related forest decline and the current consequences of historical land uses.

Although the MTEs are the focus on our evaluation and are the direct targets of our recommendations, we believe that these ecosystems serve as good reference laboratories for global change research more broadly and that our broad recommendations can be applied to other ecosystem types. Global change research in MTEs, and especially in the Mediterranean Basin, is mature enough to move a step forward and launch into an integrative phase. In this phase, research on global issues should become more inclusive and allow the development of joint projections on how ecosystems and the services they provide are expected to react to different scenarios of future change. This is the information that societies are likely to need in order to adapt to the new, uncertain changes to come.

ACKNOWLEDGEMENTS

The present review is an outcome of the research project MONTES-Consolider (CSD2008-00040), funded by the Spanish Ministry of Economy and Competitiveness, which also partially supported L.L. through the grant Complexity-NET RESINEE. X.P. is the recipient of an ICREA Acadèmia Excellence in Research grant (2011–2015). We would like to thank Joan Pino, Eduard Pla and two anonymous referees for their help and advice.

REFERENCES

- Ackerly, D.D. (2006) Plant evolution at the crossroads: the imprint of history, climate, and humans on the Mediterranean flora. *American Journal of Botany*, **93**, 170–174.
- Adams, H.D., Luce, C.H., Breshears, D.D., Allen, C.D., Weiler, M., Hale, V.C., Smith, A.M.S. & Huxman, T.E. (2012) Ecohydrological consequences of drought- and infestation-triggered tree die-off: insights and hypotheses. *Ecohydrology*, **5**, 145–159.
- Anderegg, W.R.L., Kane, J.M. & Anderegg, L.D.L. (2013) Consequences of widespread tree mortality triggered by drought and temperature stress. *Nature Climate Change*, **3**, 30–36.
- Aparicio, A., Albaladejo, R.G., Olalla-Tárraga, M.A., Carrillo, L.F. & Rodríguez, M.A. (2008) Dispersal potentials determine responses of woody plant species richness to environmental factors in fragmented Mediterranean landscapes. *Forest Ecology and Management*, **255**, 2894–2906.
- Asensio, D., Curiel Yuste, J., Mattana, S., Ribas, A., Llusà, J. & Peñuelas, J. (2012) Litter VOCs induce changes in soil microbial biomass C and N and largely increase soil CO₂ efflux. *Plant and Soil*, **360**, 163–174.
- Baldocchi, D.D., Xu, L. & Kiang, N. (2004) How plant functional-type, weather, seasonal drought, and soil physical properties alter water and energy fluxes of an oak–grass savanna and an annual grassland. *Agricultural and Forest Meteorology*, **123**, 13–39.
- Baldocchi, D.D., Ma, S., Rambal, S., Misson, L., Ourcival, J.-M., Limousin, J.-M., Pereira, J. & Papale, D. (2010) On the differential advantages of evergreenness and deciduousness in Mediterranean oak woodlands: a flux perspective. *Ecological Applications*, **20**, 1583–1597.
- Balser, T.C. & Wixon, D.L. (2009) Investigating biological control over soil carbon temperature sensitivity. *Global Change Biology*, **15**, 2935–2949.
- Barbeta, A., Ogaya, R. & Peñuelas, J. (2013) Dampening effects of long-term experimental drought on growth and mortality rates of a holm oak forest. *Global Change Biology*, **19**, 3133–3144.
- Batllori, E., Parisien, M.A., Krawchuk, M.A. & Moritz, M.A. (2013) Climate change-induced shifts in fire for Mediterranean ecosystems. *Global Ecology and Biogeography*, **22**, 1118–1129.
- Beier, C., Beierkuhnlein, C., Wohlgemuth, T., Peñuelas, J., Emmett, B., Körner, C., de Boeck, H., Christensen, J.H., Leuzinger, S., Janssens, I.A. & Hansen, K. (2012) Precipitation manipulation experiments – challenges and recommendations for the future. *Ecology Letters*, **15**, 899–911.
- Bell, T. & Adams, M. (2008) Smoke from wildfires and prescribed burning in Australia: effects on human health and ecosystems. *Wildland fires and air pollution*. (ed. by A. Bytnerowicz, M.J. Arbaugh, A.R. Riebau and C. Andersen), pp. 289–316. *Developments in environmental science* 8. Elsevier, The Hague.
- Berberoglu, S. & Akin, A. (2009) Assessing different remote sensing techniques to detect land use/cover changes in the eastern Mediterranean. *International Journal of Applied Earth Observation and Geoinformation*, **11**, 46–53.
- Beven, K. (2011) I believe in climate change but how precautionary do we need to be in planning for the future? *Hydrological Processes*, **25**, 1517–1520.
- Boer, M.M., Sadler, R.J., Wittkuhn, R.S., McCaw, L. & Grierson, P.F. (2009) Long-term impacts of prescribed burning on regional extent and incidence of wildfires – evidence from 50 years of active fire management in SW Australian forests. *Forest Ecology and Management*, **259**, 132–142.
- Boix-Fayos, C., Calvo-Cases, A., Imeson, A.C., Soriano-Soto, M.D. & Tiemessen, I.R. (1998) Spatial and short-term temporal variations in runoff, soil aggregation and other soil properties along a Mediterranean climatological gradient. *Catena*, **33**, 123–128.
- Bonan, G.B. (2008) Forests and climate change: forcings, feedbacks, and the climate benefits of forests. *Science*, **320**, 1444–1449.
- Bonet, F.J., Pérez-Luque, A.J., Moreno-Llorca, R. & Zamora, R. (2010) *Sierra Nevada global change observatory – structure and basic data*. Consejería de Medio Ambiente, Junta de Andalucía, Granada.
- Botes, A., McGeoch, M.A., Robertson, H.G., van Niekerk, A., Davids, H.P. & Chown, S.L. (2006) Ants, altitude and change

- in the northern Cape Floristic Region. *Journal of Biogeography*, **33**, 71–90.
- Brauch, H.G. (2003) Towards a fourth phase of research on human and environmental security and peace: conceptual conclusions. *Security and the environment in the Mediterranean: conceptualising security and environmental conflicts* (ed. by H.G. Brauch, P.H. Liotta, A. Marquina, P.F. Rogers and M.E.-S. Selim), pp. 919–953. Springer, Berlin.
- Brotos, L., Herrando, S. & Martín, J.L. (2004) Bird assemblages in forest fragments within Mediterranean mosaics created by wild fires. *Landscape Ecology*, **19**, 663–675.
- Brotos, L., Aquilué, N., de Cáceres, M., Fortin, M.-J. & Fall, A. (2013) How fire history, fire suppression practices and climate change affect wildfire regimes in Mediterranean landscapes. *PLoS ONE*, **8**, e62392.
- Brown, S., Mo, J.M., McPherson, J.K. & Bell, D.T. (1996) Decomposition of woody debris in Western Australian forests. *Canadian Journal of Forest Research*, **26**, 954–966.
- Campos, P., Huntsinger, L., Oviedo, J.L., Díaz, M., Starrs, P., Standiford, R.B. & Montero, G. (2013) *Mediterranean oak woodland working landscapes: Dehesas of Spain and Ranchlands of California*. Springer, New York.
- Carnicer, J., Coll, M., Ninyerola, M., Pons, X., Sánchez, G. & Peñuelas, J. (2011) Widespread crown condition decline, food web disruption, and amplified tree mortality with increased climate change-type drought. *Proceedings of the National Academy of Sciences USA*, **108**, 1474–1478.
- Castro, J., Allen, C.D., Molina-Morales, M., Marañón-Jiménez, S., Sánchez-Miranda, A. & Zamora, R. (2011) Salvage logging versus the use of burnt wood as a nurse object to promote post-fire tree seedling establishment. *Restoration Ecology*, **19**, 537–544.
- Castro, J., Puerta-Piñero, C., Leverkus, A.B., Moreno-Rueda, G. & Sánchez-Miranda, A. (2012) Post-fire salvage logging alters a key plant–animal interaction for forest regeneration. *Ecosphere*, **3**(10), art. 90. doi: 10.1890/ES12-00089
- Cheai, A., Badeau, V., Boe, J., Chuine, I., Delire, C., Dufréne, E., François, C., Gritti, E.S., Legay, M., Pagé, C., Thuiller, W., Viomy, N. & Leadley, P. (2012) Climate change impacts on tree ranges: model intercomparison facilitates understanding and quantification of uncertainty. *Ecology Letters*, **15**, 533–544.
- Ciccioli, P., Brancaleoni, E., Frattoni, M., Cecinato, A. & Pinciarelli, L. (2001) Determination of volatile organic compounds (VOC) emitted from biomass burning of Mediterranean vegetation species by GC-MS. *Analytical Letters*, **34**, 937–955.
- Clavero, M., Villero, D. & Brotos, L. (2011) Climate change or land use dynamics: do we know what climate change indicators indicate? *PLoS ONE*, **6**, e18581.
- Conacher, A.J. (1998) Summary and conclusions. *Land degradation in Mediterranean environments of the world; nature and extent, causes and solutions* (ed. by A.J. Conacher and M. Sala), pp. 449–457. John Wiley and Sons, Chichester.
- Concepción, E.D., Díaz, M. & Baquero, R.A. (2008) Effects of landscape complexity on the ecological effectiveness of agri-environment schemes. *Landscape Ecology*, **23**, 135–148.
- Concepción, E.D., Díaz, M., Kleijn, D., Báldi, A., Batáry, P., Clough, Y., Gabriel, D., Herzog, F., Holzschuh, A., Knop, E., Marshall, J.P., Tschardtke, T. & Verhulst, J. (2012) Interactive effects of landscape context constrains the effectiveness of local agri-environmental management. *Journal of Applied Ecology*, **49**, 695–705.
- Crecente-Campo, F., Pommerening, A. & Rodríguez-Soalleiro, R. (2009) Impacts of thinning on structure, growth and risk of crown fire in a *Pinus sylvestris* L. plantation in northern Spain. *Forest Ecology and Management*, **257**, 1945–1954.
- Curiel Yuste, J., Baldocchi, D.D., Gershenson, A., Goldstein, A., Misson, L. & Wong, S. (2007) Microbial soil respiration and its dependency on carbon inputs, soil temperature and moisture. *Global Change Biology*, **13**, 2018–2035.
- Curiel Yuste, J., Barba, J., Fernandez-Gonzalez, M., Fernandez-Lopez, A.J., Mattana, S., Martinez-Vilalta, J., Nolis, P. & Lloret, F. (2012) Changes in soil bacterial community triggered by drought-induced gap succession preceded changes in soil C stocks and quality. *Ecology and Evolution*, **2**, 3016–3031.
- D'Agostino, D.R., Trisorio, L.G., Lamaddalena, N. & Ragab, R. (2010) Assessing the results of scenarios of climate and land use changes on the hydrology of an Italian catchment: modelling study. *Hydrological Processes*, **24**, 2693–2704.
- Delgado, J., Llorens, P., Nord, G., Calder, I.R. & Gallart, F. (2010) Modelling the hydrological response of a Mediterranean medium-sized headwater basin subject to land cover change: the Cardener River basin (NE Spain). *Journal of Hydrology*, **383**, 125–134.
- Díaz, M., Tietje, W. & Barret, R. (2013) Effects of management on biological diversity and endangered species. *Mediterranean oak woodland working landscapes: Dehesas of Spain and Ranchlands of California* (ed. by P. Campos, L. Huntsinger, J.L. Oviedo, M. Díaz, P. Starrs, R.B. Standiford and G. Montero), pp. 213–243. Springer, New York.
- Díaz-Delgado, R. & Pons, X. (2001) Spatial patterns of forest fires in Catalonia (NE of Spain) along the period 1975–1995. Analysis of vegetation recovery after fire. *Forest Ecology and Management*, **147**, 67–74.
- Díaz-Delgado, R., Lloret, F., Pons, X. & Terradas, J. (2002) Satellite evidence of decreasing resilience in Mediterranean plant communities after recurrent wildfires. *Ecology*, **83**, 2293–2303.
- Dimitrakopoulos, P.G. (2010) Influence of evenness on the litter-species richness–decomposition relationship in Mediterranean grasslands. *Journal of Plant Ecology*, **3**, 71–78.
- Dimitrakopoulos, P.G., Siamantziouras, A.-S.D., Galanidis, A., Mprezetou, I. & Troumbis, A.Y. (2006) The interactive effects of fire and diversity on short-term responses of ecosystem processes in experimental Mediterranean grasslands. *Environmental Management*, **37**, 826–839.
- Doblas-Miranda, E., Sánchez-Piñero, F. & González-Megías, A. (2009) Different structuring factors but connected dynamics shape litter and belowground soil macrofaunal food webs. *Soil Biology and Biochemistry*, **41**, 2543–2550.

- Dunford, R., Michel, K., Gagnage, M., Piégay, H. & Trémel, M.L. (2009) Potential and constraints of unmanned aerial vehicle technology for the characterization of Mediterranean riparian forest. *International Journal of Remote Sensing*, **30**, 4915–4935.
- Eagleson, P.S. (1982) Ecological optimality in water-limited natural soil–vegetation systems: 1. Theory and hypothesis. *Water Resources Research*, **18**, 325–340.
- Emmett, B.A., Beier, C., Estiarte, M., Tietema, A., Kristensen, H.L., Williams, D., Peñuelas, J., Schmidt, I. & Sowerby, A. (2004) The response of soil processes to climate change: results from manipulation studies of shrublands across an environmental gradient. *Ecosystems*, **7**, 625–637.
- Estornell, J., Ruiz, L.A. & Velázquez-Martí, B. (2010) Study of shrub cover and height using LIDAR data in a Mediterranean area. *Forest Science*, **57**, 171–179.
- Ewen, J., O'Donnell, G., Burton, A. & O'Connell, E. (2006) Errors and uncertainty in physically-based rainfall-runoff modelling of catchment change effects. *Journal of Hydrology*, **330**, 641–650.
- Fatichi, S., Leuzinger, S. & Körner, C. (2014) Moving beyond photosynthesis: from carbon source to sink-driven vegetation modeling. *New Phytologist*, **201**, 1086–1095.
- Fernandes, P.M., Rego, F.C. & Rigolot, E. (2011) The FIRE PARADOX project: towards science-based fire management in Europe. *Forest Ecology and Management*, **261**, 2177–2178.
- Franklin, J. (2010) Vegetation dynamics and exotic plant invasion following high severity crown fire in a southern California conifer forest. *Plant Ecology*, **207**, 281–295.
- Franklin, J., Syphard, A.D., He, H.S. & Mladenoff, D.J. (2005) Altered fire regimes affect landscape patterns of plant succession in the foothills and mountains of southern California. *Ecosystems*, **8**, 885–898.
- Garbulsky, M.F., Peñuelas, J., Papale, D. & Filella, I. (2008) Remote estimation of carbon dioxide uptake by a Mediterranean forest. *Global Change Biology*, **14**, 2860–2867.
- García, M., Riaño, D., Chuvieco, E. & Dansond, F.M. (2010) Estimating biomass carbon stocks for a Mediterranean forest in central Spain using LiDAR height and intensity data. *Remote Sensing of Environment*, **114**, 816–830.
- García, D., Zamora, R. & Amico, G.C. (2011) The spatial scale of plant–animal interactions: effects of resource availability and habitat structure. *Ecological Monographs*, **81**, 103–121.
- García-Hurtado, E., Pey, J., Baeza, M.J., Carrara, A., Llovet, J., Querol, X., Alastuey, A. & Vallejo, V.R. (2013) Carbon emissions in Mediterranean shrubland wildfires: an experimental approach. *Atmospheric Environment*, **69**, 86–93.
- García-Palacios, P., Bowker, M.A., Chapman, S.J., Maestre, F.T., Soliveres, S., Gallardo, A., Valladares, F., Guerrero, C. & Escudero, A. (2011) Early-successional vegetation changes after roadside prairie restoration modify processes related with soil functioning by changing microbial functional diversity. *Soil Biology and Biochemistry*, **43**, 1245–1253.
- Garnier, E., Lavorel, S., Ansquer, P. *et al.* (2007) Assessing the effects of land-use change on plant traits, communities and ecosystem functioning in grasslands: a standardized methodology and lessons from an application to 11 European sites. *Annals of Botany*, **99**, 967–985.
- Gil-Tena, A., Brotons, L. & Saura, S. (2009) Mediterranean forest dynamics and forest bird distribution changes in the late 20th century. *Global Change Biology*, **15**, 474–485.
- Giorgi, F. (2006) Climate change hot-spots. *Geophysical Research Letters*, **33**, L08707.
- Giorgi, F. & Lionello, P. (2008) Climate change projections for the Mediterranean region. *Global and Planetary Change*, **63**, 90–104.
- Glenn, E.P., Doody, T.M., Guerschman, J.P., Huete, A.R., King, E.A., McVicar, T.R., Van Dijk, A.I.J.M., Van Niel, T.G., Yebra, M. & Zhang, Y. (2011) Actual evapotranspiration estimation by ground and remote sensing methods: the Australian experience. *Hydrological Processes*, **26**, 4103–4116.
- Gómez-Aparicio, L., Zamora, R., Gómez, J.M., Hódar, J.A., Castro, J. & Baraza, E. (2004) Applying plant facilitation to forest restoration: a meta-analysis of the use of shrubs as nurse plants. *Ecological Applications*, **14**, 1128–1138.
- Gonçalves, C., Alves, C., Fernandes, A.P., Monteiro, C., Tarelho, L., Evtyugina, M. & Pio, C. (2011) Organic compounds in PM_{2.5} emitted from fireplace and woodstove combustion of typical Portuguese wood species. *Atmospheric Environment*, **45**, 4533–4545.
- Granier, A., Reichstein, M., Breda, N. *et al.* (2007) Evidence for soil water control on carbon and water dynamics in European forests during the extremely dry year: 2003. *Agricultural and Forest Meteorology*, **143**, 123–145.
- Grigulis, K., Lavorel, S., Davies, I.D., Dossantos, A., Lloret, F. & Vila, M. (2005) Landscape-scale positive feedbacks between fire and expansion of the large tussock grass, *Ampelodesmos mauritanica* in Catalan shrublands. *Global Change Biology*, **11**, 1042–1053.
- Gritti, E.S., Smith, B. & Sykes, M.T. (2006) Vulnerability of Mediterranean Basin ecosystems to climate change and invasion by exotic plant species. *Journal of Biogeography*, **33**, 145–157.
- Heck, P., Lfithi, D., Wernli, H. & Schar, C. (2001) Climate impacts of European scale anthropogenic vegetation changes: a sensitivity study using a regional climate model. *Journal of Geophysical Research*, **106**, 7817–7835.
- Hernández-Clemente, R., Navarro-Cerrillo, R.M. & Zarco-Tejada, P.J. (2012) Carotenoid content estimation in a heterogeneous conifer forest using narrow-band indices and PROSPECT plus DART simulations. *Remote Sensing of Environment*, **127**, 298–315.
- Hernández-Santana, V., Martínez-Vilalta, J., Martínez-Fernández, J. & Williams, M. (2009) Evaluating the effect of drier and warmer conditions on water use by *Quercus pyrenaica*. *Forest Ecology and Management*, **258**, 1719–1730.
- Hoff, C. & Rambal, S. (2003) An examination of the interaction between climate, soil and leaf area index in a *Quercus ilex* ecosystem. *Annals of Forest Science*, **60**, 153–161.
- Jacobsen, A.L., Pratt, R.B., Ewers, F.W. & Davis, S.D. (2007) Cavitation resistance among 26 chaparral species of southern California. *Ecological Monographs*, **77**, 99–115.

- Janion, C., Bedos, A., Bengtsson, J., Deharveng, L., van Vuuren, B.J., Leinaas, H.P., Liu, A., Malmstrom, A., Porcos, D. & Chown, S.L. (2011) Springtail diversity in South Africa. *South African Journal of Science*, **107**, 75–81.
- Jump, A.S., Marchant, R. & Peñuelas, J. (2008) Environmental change and the option value of genetic diversity. *Trends in Plant Science*, **14**, 51–58.
- Keane, R.E., Cary, G.J., Davies, I.D., Flannigan, M.D., Gardner, R.H., Lavorel, S., Lenihan, J.M., Li, C. & Rupp, T.S. (2004) A classification of landscape fire succession models: spatial simulations of fire and vegetation dynamics. *Ecological Modelling*, **179**, 3–27.
- Keeley, J.E. & Brennan, T.J. (2012) Fire-driven alien invasion in a fire-adapted ecosystem. *Oecologia*, **169**, 1043–1052.
- Keeley, J.E. & Zedler, P.H. (2009) Large, high-intensity fire events in southern California shrublands: debunking the fine-grain age patch model. *Ecological Applications*, **19**, 69–94.
- Keeley, J.E., Bond, W.J., Bradstock, R.A., Pausas, J.G. & Rundel, P.W. (2012) *Fire in Mediterranean ecosystems: ecology, evolution and management*. Cambridge University Press, Cambridge.
- Keenan, T., Sabate, S. & Gracia, C. (2010) Soil water stress and coupled photosynthesis–conductance models: bridging the gap between conflicting reports on the relative roles of stomatal, mesophyll conductance and biochemical limitations to photosynthesis. *Agricultural and Forest Meteorology*, **150**, 443–453.
- Keenan, T.F., Serra, P., Lloret, F., Ninyerola, M. & Sabate, S. (2011) Predicting the future of forests in the Mediterranean under climate change, with niche- and process-based models: CO₂ matters! *Global Change Biology*, **17**, 265–279.
- Kerkhoff, A.J., Martens, S.N. & Milne, B.T. (2004) An ecological evaluation of Eagleson's optimality hypotheses. *Functional Ecology*, **18**, 404–413.
- Kraaij, T., Cowling, R.M. & van Wilgen, B.W. (2013) Lightning and fire weather in eastern coastal fynbos shrublands: seasonality and long-term trends. *International Journal of Wildland Fire*, **22**, 288–295.
- Kurt, Y., González-Martínez, S.C., Alia, R. & Isik, K. (2012) Genetic differentiation in *Pinus brutia* Ten. using molecular markers and quantitative traits: the role of altitude. *Annals of Forest Science*, **69**, 345–351.
- Larsen, T.H., Williams, N.M. & Kremen, C. (2005) Extinction order and altered community structure rapidly disrupt ecosystem functioning. *Ecology Letters*, **8**, 538–547.
- Lau, J.A. & Lennon, J.T. (2012) Rapid responses of soil microorganisms improve plant fitness in novel environments. *Proceedings of the National Academy of Sciences USA*, **109**, 14058–14062.
- Lavorel, S. (1999) Ecological diversity and resilience of Mediterranean vegetation to disturbance. *Diversity and Distributions*, **5**, 1–2.
- Lavorel, S. & Garnier, E. (2002) Predicting changes in community composition and ecosystem functioning from plant traits: revisiting the Holy Grail. *Functional Ecology*, **16**, 545–556.
- Lavorel, S., Canadell, J., Rambal, S. & Terradas, J. (1998) Mediterranean terrestrial ecosystems: research priorities on global change effects. *Global Ecology and Biogeography Letters*, **7**, 157–166.
- Lazaridis, M., Spyridaki, A., Solberg, S., Smolik, J., Zdimal, V., Eleftheriadis, K., Aleksandropoulou, V., Hov, Ø. & Georgopoulos, P.G. (2005) Mesoscale modelling of combined aerosol and photooxidant processes in the eastern Mediterranean. *Atmospheric Chemistry and Physics*, **5**, 927–940.
- Le Page, Y., Oom, D., Silva, J.M.N., Jönsson, P. & Pereira, J.M.C. (2010) Seasonality of vegetation fires as modified by human action: observing the deviation from eco-climatic fire regimes. *Global Ecology and Biogeography*, **19**, 575–588.
- Lee, D.E., Bond, M.L., Borchert, M.I. & Tanner, R. (2013) Influence of fire and salvage logging on site occupancy of spotted owls in the San Bernardino and San Jacinto Mountains of southern California. *Journal of Wildlife Management*, **77**, 1327–1341.
- Lenihan, J.M., Drapek, R., Bachelet, D. & Neilson, R.P. (2003) Climate change effects on vegetation distribution, carbon, and fire in California. *Ecological Applications*, **13**, 1667–1681.
- Lespinas, F., Ludwig, W. & Heussner, S. (2010) Impact of recent climate change on the hydrology of coastal mediterranean rivers in southern France. *Climatic Change*, **99**, 425–456.
- Leverkus, A.B., Lorite, J., Navarro, F.B., Sánchez-Cañete, E.P. & Castro, J. (2013) Post-fire salvage logging alters species composition and reduces cover, richness, and diversity in Mediterranean plant communities. *Journal of Environmental Management*, **133**, 323–331.
- Li, J.-H., Dugas, W.A., Hymus, G.J., Johnson, D.P., Hinkle, C.R., Drake, B.G. & Hungate, B.A. (2003) Direct and indirect effects of elevated CO₂ on transpiration from *Quercus myrtifolia* in a scrub-oak ecosystem. *Global Change Biology*, **9**, 96–105.
- Limousin, J.M., Rambal, S., Ourcival, J.M., Rocheteau, A., Joffre, R. & Rodriguez-Cortina, R. (2009) Long-term transpiration change with rainfall decline in a Mediterranean *Quercus ilex* forest. *Global Change Biology*, **15**, 2163–2175.
- Lionello, P., Malanotte-Rizzoli, P. & Boscolo, R. (2006) *Mediterranean climate variability*. Elsevier, Amsterdam.
- Liu, Y., Stanturf, J. & Goodrick, S. (2010) Trends in global wild-fire potential in a changing climate. *Forest Ecology and Management*, **259**, 685–697.
- Lloret, F., Calvo, E., Pons, X. & Diaz-Delgado, R. (2002) Wildfires and landscape patterns in the eastern Iberian Peninsula. *Landscape Ecology*, **17**, 745–759.
- Lloret, F., Estevan, H., Vayreda, J. & Terradas, J. (2005) Fire regenerative syndromes of forest woody species across fire and climatic gradients. *Oecologia*, **146**, 461–468.
- Lloret, F., Peñuelas, J., Prieto, P., Llorens, L. & Estiarte, M. (2009) Plant community changes induced by experimental climate change: seedling and adult species composition. *Perspectives in Plant Ecology, Evolution and Systematic*, **11**, 53–63.
- Lloret, F., Escudero, A., Iriondo, J.M., Martínez-Vilalta, J. & Valladares, F. (2012) Extreme climatic events and vegetation: the role of stabilizing processes. *Global Change Biology*, **18**, 797–805.

- Llovet, J., Ruiz-Valera, M., Josa, R. & Vallejo, V.R. (2009) Soil responses to fire in Mediterranean forest landscapes in relation to the previous stage of land abandonment. *International Journal of Wildland Fire*, **18**, 222–232.
- Loepfe, L., Martínez-Vilalta, J. & Piñol, J. (2011) An integrative model of human-influenced fire regimes and landscape dynamics. *Environmental Modelling and Software*, **26**, 1028–1040.
- Loepfe, L., Rodrigo, A. & Lloret, F. (2012) Two thresholds determine climatic control of forest-fire size in Europe. *Biogeosciences Discussions*, **9**, 9065–9089.
- López-Moreno, J.I., Vicente-Serrano, S.M., Moran-Tejeda, E., Zabalza, J., Lorenzo-Lacruz, J. & García-Ruiz, J.M. (2011) Impact of climate evolution and land use changes on water yield in the Ebro Basin. *Hydrology and Earth System Sciences*, **15**, 311–322.
- Lu, D. (2006) The potential and challenge of remote sensing-based biomass estimation. *International Journal of Remote Sensing*, **27**, 1297–1328.
- McDowell, N.G., Ryan, M.G., Zeppel, M.J.B. & Tissue, D.T. (2013) Feature: improving our knowledge of drought-induced forest mortality through experiments, observations, and modeling. *New Phytologist*, **200**, 289–293.
- Macfarlane, C., Lardner, T., Patterson, K. & Grigg, A.H. (2010) A new model for predicting understorey leaf area from biomass in eucalypt forest to test the ecohydrological equilibrium theory. *Methods in Ecology and Evolution*, **1**, 371–379.
- Maestre, F.T., Valladares, F. & Reynolds, J.F. (2005) Is the change of plant–plant interactions with abiotic stress predictable? A meta-analysis of field results in arid environments. *Journal of Ecology*, **93**, 748–757.
- Maestre, F.T., Castillo-Monroy, A.P., Bowker, M.A. & Ochoa-Hueso, R. (2012) Species richness effects on ecosystem multifunctionality depend on evenness, composition and spatial pattern. *Journal of Ecology*, **100**, 317–330.
- Malcolm, J.R., Liu, C., Neilson, R.P., Hansen, L. & Hannah, L. (2006) Global warming and extinctions of endemic species from biodiversity hotspots. *Conservation Biology*, **20**, 538–548.
- Marañón-Jiménez, S. & Castro, J. (2013) Effect of decomposing post-fire coarse woody debris on soil fertility and nutrient availability in a Mediterranean ecosystem. *Biogeochemistry*, **112**, 519–535.
- Martínez-Vilalta, J., Lloret, F. & Breshears, D.D. (2012) Drought-induced forest decline: causes, scope and implications. *Biology Letters*, **8**, 689–691.
- Marzano, R., Garbarino, M., Marcolin, E., Pividori, M. & Lingua, E. (2013) Deadwood anisotropic facilitation on seedling establishment after a stand-replacing wildfire in Aosta Valley (NW Italy). *Ecological Engineering*, **51**, 117–122.
- Matias, L., Quero, J.L., Zamora, R. & Castro, J. (2012) Evidence for plant traits driving specific drought resistance. A community field experiment. *Environmental Experimental Botany*, **81**, 55–61.
- Millington, J.D., Wainwright, J., Perry, G.L.W., Romero-Calcerrada, R.R. & Malamud, B.D. (2009) Modelling Mediterranean landscape succession-disturbance dynamics: a landscape fire-succession model. *Environmental Modelling and Software*, **24**, 1196–1208.
- Misson, L., Degueldre, D., Collin, C., Rodríguez, R., Rocheteau, A., Ourcival, J.-M. & Rambal, S. (2011) Phenological responses to extreme droughts in a Mediterranean forest. *Global Change Biology*, **17**, 1036–1048.
- Mitsopoulos, I.D. & Dimitrakopoulos, A.P. (2007) Canopy fuel characteristics and potential crown fire behavior in Aleppo pine (*Pinus halepensis* Mill.) forests. *Annals of Forest Science*, **64**, 287–299.
- Monson, R.K., Trahan, N., Rosenstiel, T.N., Veres, P., Moore, D., Wilkinson, M., Norby, R.J., Volder, A., Tjoelker, M.G., Briske, D.D., Karnosky, D.F. & Fall, R. (2007) Isoprene emission from terrestrial ecosystems in response to global change: minding the gap between models and observations. *Philosophical Transactions of the Royal Society A*, **365**, 1677–1695.
- Montenegro, G., Ginocchio, R., Segura, A., Keeley, J.E. & Gomez, M. (2004) Fire regimes and vegetation responses in two Mediterranean-climate regions. *Revista Chilena de Historia Natural*, **77**, 455–464.
- Morán-Tejeda, E., López-Moreno, J.I., Ceballos-Barbancho, A. & Vicente-Serrano, S.M. (2011) River regimes and recent hydrological changes in the Duero Basin (Spain). *Journal of Hydrology*, **404**, 241–258.
- Moreira, F., Arianoutsou, M., Corona, P. & De Las Heras, J. (2012) *Post-fire management and restoration of southern European forests*. Springer, Berlin.
- Moreira, F., Ferreira, A., Abrantes, N., Catry, F., Fernandes, P., Roxo, L., Keizer, J.J. & Silva, J. (2013) Occurrence of native and exotic invasive trees in burned pine and eucalypt plantations: implications for post-fire forest conversion. *Ecological Engineering*, **58**, 296–302.
- Mouillot, F., Rambal, S. & Joffre, R. (2002) Simulating climate change impacts on fire frequency and vegetation dynamics in a Mediterranean-type ecosystem. *Global Change Biology*, **8**, 423–437.
- Muhle, J., Lueker, T.J., Su, Y., Miller, B.R., Prather, K.A. & Weiss, R.F. (2007) Trace gas and particulate emissions from the 2003 southern California wildfires. *Journal of Geophysical Research*, **112**, D03307.
- Munné, A. & Prat, N. (2011) Effects of Mediterranean climate annual variability on stream biological quality assessment using macroinvertebrate communities. *Ecological Indicators*, **11**, 651–662.
- Navarro-González, I., Pérez-Luque, A.J., Bonet, F.J. & Zamora, R. (2013) The weight of the past: land-use legacies and recolonization of pine plantations by oak trees. *Ecological Applications*, **23**, 1267–1276.
- Ortega, M., Gómez-Sanz, V., García del Barrio, J.M. & Elena Roselló, R. (2010) *Monitoring vulnerability of the Spanish forest landscapes: the SISPAES approach*. IUFRO Landscape Ecology Working Group Conference, Bragança.
- Otero, I., Boada, M., Badia, A., Pla, E., Vayreda, J., Sabaté, S., Gracia, C.A. & Peñuelas, J. (2011) Loss of water availability and stream biodiversity under land abandonment and climate

- change in a Mediterranean catchment (Olzinelles, NE Spain). *Land Use Policy*, **28**, 207–218.
- Pasquini, S.C. & Vourlitis, G.L. (2010) Post-fire primary production and plant community dynamics in chaparral stands exposed to varying levels of nitrogen deposition. *Journal of Arid Environments*, **74**, 310–314.
- Paton-Walsh, C., Louisa, K., Emmons, L.K. & Wiedinmyer, C. (2012) Australia's Black Saturday fires – comparison of techniques for estimating emissions from vegetation fires. *Atmospheric Environment*, **60**, 262–270.
- Pausas, J.G. (1999) Response of plant functional types to changes in the fire regime in Mediterranean ecosystems: a simulation approach. *Journal of Vegetation Science*, **10**, 717–722.
- Pausas, J.G. (2004) Changes in fire and climate in the eastern Iberian Peninsula (Mediterranean Basin). *Climate Change*, **63**, 337–350.
- Peñuelas, J. & Staudt, M. (2010) BVOCs and global change. *Trends in Plant Science*, **15**, 133–144.
- Peñuelas, J., Guenther, A., Rapparini, F. *et al.* (2013) Intensive measurements of gas, water, and energy exchange between vegetation and troposphere during the MONTES campaign in a vegetation gradient from short semi-desertic shrublands to tall wet temperate forests in the NW Mediterranean Basin. *Atmospheric Environment*, **75**, 348–364.
- Pérez, B., Cruz, A., Fernández-González, F. & Moreno, J.M. (2003) Effects of the recent land-use history on the postfire vegetation of uplands in central Spain. *Forest Ecology and Management*, **182**, 273–283.
- Peterman, W., Waring, R.H., Seager, T. & Pollock, W.L. (2013) Soil properties affect pinyon pine–juniper response to drought. *Ecohydrology*, **6**, 455–463.
- Petit, R.J., Hampe, A. & Cheddadi, R. (2005) Climate changes and tree phylogeography in the Mediterranean. *Taxon*, **54**, 877–885.
- Petit, R.J., Hu, F.S. & Dick, C.W. (2008) Forests of the past: a window to future changes. *Science*, **320**, 1450–1452.
- Phuleria, H.C., Fine, P.M., Zhu, Y. & Sioutas, C. (2005) Air quality impacts of the October 2003 Southern California wildfires. *Journal of Geophysical Research*, **110**, D07S20.
- Pino, J., Arnan, X., Rodrigo, A. & Retana, J. (2013) Post-fire invasion and subsequent extinction of *Conyza* spp. in Mediterranean forests is mostly explained by local factors. *Weed Research*, **53**, 470–478.
- Piñol, J., Castellnou, M. & Beven, K.J. (2007) Conditioning uncertainty in ecological models: assessing the impact of fire management strategies. *Ecological Modelling*, **207**, 34–44.
- Pitman, A.J., Narisma, G.T., Pielke, R.A. Sr & Holbrook, N.J. (2004) Impact of land cover change on the climate of southwest Western Australia. *Journal of Geophysical Research*, **109**, D18109.
- Prentice, I.C., Kelley, D.I., Foster, P.N., Friedlingstein, P., Harrison, S.P. & Bartlein, P.J. (2011) Modeling fire and the terrestrial carbon balance. *Global Biogeochemical Cycles*, **25**, GB3005.
- Price, O.F. & Bradstock, R.A. (2010) The effect of fuel age on the spread of fire in a sclerophyll forest in the Sydney region of Australia. *International Journal of Wildland Fire*, **19**, 35–45.
- Prieur-Richard, A.-H. & Lavorel, S. (2000) Invasions: the perspective of diverse plant communities – a review. *Australian Ecology*, **25**, 1–7.
- Prieur-Richard, A.-H., Lavorel, S., Linhart, Y.B. & Dos Santos, A. (2002) Plant diversity, herbivory and resistance of a plant community to invasion in Mediterranean annual communities. *Oecologia*, **130**, 96–104.
- Puerta-Piñero, C., Espelta, J.M., Sánchez-Humanes, B., Rodrigo, A., Coll, L. & Brotons, L. (2012) History matters: previous land use changes determine post-fire vegetation recovery in forested Mediterranean landscapes. *Forest Ecology and Management*, **279**, 121–127.
- Quero, J., Sterck, F., Martínez-Vilalta, J. & Villar, R. (2011) Water-use strategies of six co-existing Mediterranean woody species during a summer drought. *Oecologia*, **166**, 45–57.
- Ramírez-Valiente, J.A., Lorenzo, Z., Soto, A., Valladares, F., Gil, L. & Aranda, I. (2009) Elucidating the role of genetic drift and natural selection in cork oak differentiation regarding drought tolerance. *Molecular Ecology*, **18**, 3803–3815.
- Raz-Yaseef, N., Yakir, D., Schiller, G. & Cohen, S. (2012) Dynamics of evapotranspiration partitioning in a semi-arid forest as affected by temporal rainfall patterns. *Agricultural and Forest Meteorology*, **157**, 77–85.
- Reichstein, M., Tenhunen, J., Rouspard, O., Ourcival, J.-M., Rambal, S., Miglietta, F., Peressotti, A., Pecchiari, M., Tirone, G. & Valentini, R. (2003) Inverse modeling of seasonal drought effects on canopy CO₂/H₂O exchange in three Mediterranean ecosystems. *Journal of Geophysical Research*, **108**, 4726.
- Rey Benayas, J.M., Martins, A., Nicolau, J.M. & Schulz, J.J. (2007) Abandonment of agricultural land: an overview of drivers and consequences. *CAB Reviews: Perspectives in Agriculture, Veterinary Science, Nutrition and Natural Resources*, **2**, 057, doi: 10.1079/PAVSNNR20072057.
- Rouget, M., Richardson, D.M., Milton, S.J. & Polakow, D. (2001) Predicting invasion dynamics of four alien *Pinus* species in a highly fragmented semi-arid shrubland in South Africa. *Plant Ecology*, **152**, 79–92.
- Ruiz-Benito, P., Lines, E.R., Gómez-Aparicio, L., Zavala, M.A. & Coomes, D.A. (2013) Patterns and drivers of tree mortality in Iberian forests: climatic effects are modified by competition. *PLoS ONE*, **8**, e56843.
- Rundel, P.W. (1998) Landscape disturbance in Mediterranean-type ecosystems: an overview. *Landscape disturbance and biodiversity in Mediterranean-type ecosystems* (ed. by P.W. Rundel, G. Montenegro and F.M. Jaksic), pp. 3–22. Springer-Verlag, Berlin.
- Ruwanza, S., Gaertner, M., Esler, K.J. & Richardson, D.M. (2013) Both complete clearing and thinning of invasive trees lead to short-term recovery of native riparian vegetation in the Western Cape, South Africa. *Applied Vegetation Science*, **16**, 193–204.

- Ryals, R. & Silver, W.L. (2013) Effects of organic matter amendments on net primary productivity and greenhouse gas emissions in annual grasslands. *Ecological Applications*, **23**, 46–59.
- Sala, O.E., Chapin, F.S. III, Armesto, J.J., Berlow, E., Bloomfield, J., Dirzo, R., Huber-Sanwald, E., Huenneke, L.F., Jackson, R.B., Kinzig, A., Leemans, R., Lodge, D.M., Mooney, H.A., Oesterheld, M., Poff, N.L., Sykes, M.T., Walker, B.H., Walker, M. & Wall, D.H. (2000) Global biodiversity scenarios for the year 2100. *Science*, **287**, 1770–1774.
- Sánchez-Salguero, R., Navarro-Cerrillo, R.M., Swetnam, T.W. & Zavala, M.A. (2012) Is drought the main decline factor at the rear edge of Europe? The case of southern Iberian pine plantations. *Forest Ecology and Management*, **271**, 158–169.
- San-Miguel-Ayán, J., Moreno, J.M. & Camia, A. (2013) Analysis of large fires in European Mediterranean landscapes: lessons learned and perspectives. *Forest Ecology and Management*, **294**, 11–22.
- Santos, T., Tellería, J.L., Díaz, M. & Carbonell, R. (2006) Evaluating the environmental benefits of CAP reforms: can afforestations restore forest bird communities in Mediterranean Spain? *Basic and Applied Ecology*, **7**, 483–495.
- Sardans, J., Peñuelas, J. & Estiarte, M. (2008) Changes in soil enzymes related to C and N cycle and in soil C and N content under prolonged warming and drought in a Mediterranean shrubland. *Applied Soil Ecology*, **39**, 223–235.
- Schröter, D., Cramer, W., Leemans, R. *et al.* (2005) Ecosystem service supply and vulnerability to global change in Europe. *Science*, **310**, 1333–1337.
- Seibert, J., McDonnell, J.J. & Woodsmith, R.D. (2010) Effects of wildfire on catchment runoff response: a modelling approach to detect changes in snow-dominated forested catchments. *Hydrology Research*, **41**, 378–390.
- Selmants, P.C., Zavaleta, E.S., Pasari, J.R. & Hernandez, D.L. (2012) Realistic plant species losses reduce invasion resistance in a California serpentine grassland. *Journal of Ecology*, **100**, 723–731.
- Senatore, A., Mendicino, G., Smiatek, G. & Kunstmann, H. (2011) Regional climate change projections and hydrological impact analysis for a Mediterranean basin in southern Italy. *Journal of Hydrology*, **399**, 70–92.
- Seneviratne, S.I., Corti, T., Davin, E.L., Hirschi, M., Jaeger, E.B., Lehner, I., Orlowsky, B. & Teuling, A.J. (2010) Investigating soil moisture–climate interactions in a changing climate: a review. *Earth-Science Reviews*, **99**, 125–161.
- Serra, P., Saurí, D. & Pons, X. (2008) Land-cover and land-use in a Mediterranean landscape: a spatial analysis of driving forces integrating biophysical and human factors. *Applied Geography*, **28**, 189–209.
- Serrano, L., Penuelas, J. & Ustin, S.L. (2001) Remote sensing of nitrogen and lignin in Mediterranean vegetation from AVIRIS data: decomposing biochemical from structural signals. *Remote Sensing of Environment*, **81**, 355–364.
- Serrano-Ortiz, P., Marañón-Jiménez, S., Reverter, B.R., Sánchez-Cañete, E.P., Castro, J., Zamora, R. & Kowalski, A.S. (2011) Post-fire salvage logging reduces carbon sequestration in Mediterranean coniferous forest. *Forest Ecology and Management*, **262**, 2287–2296.
- Shakesby, R.A. (2011) Post-wildfire soil erosion in the Mediterranean: review and future research directions. *Earth-Science Reviews*, **105**, 71–100.
- Shoshany, M. (2000) Satellite remote sensing of natural Mediterranean vegetation: a review within an ecological context. *Progress in Physical Geography*, **24**, 153–178.
- Siles, G., Rey, P.J. & Alcántara, J.M. (2010) Post-fire restoration of Mediterranean forests: testing assembly rules mediated by facilitation. *Basic and Applied Ecology*, **11**, 422–431.
- Sillmann, J., Kharin, V.V., Zwiers, F.W., Zhang, X. & Bronaugh, D. (2013) Climate extremes indices in the CMIP5 multimodel ensemble: part 2. Future climate projections. *Journal of Geophysical Research: Atmospheres*, **118**, 2473–2493.
- Steffen, W., Sanderson, A., Tyson, P.D., Jäger, J., Matson, P.A., Moore, B. III, Oldfield, F., Richardson, K., Schellnhuber, H.J., Turner, B.L. & Wasson, R.J. (2004) *Global change and the earth system: a planet under pressure*. Springer-Verlag, Berlin.
- Sus, O., Williams, M., Bernhofer, C., Béziat, P., Buchmann, N., Ceschia, E., Doherty, R., Eugster, W., Grünwald, T., Kutsch, W., Smith, P. & Wattenbach, M. (2010) A linked carbon cycle and crop developmental model: description and evaluation against measurements of carbon fluxes and carbon stocks at several European agricultural sites. *Agriculture, Ecosystems and Environment*, **139**, 402–418.
- Syphard, A.D. & Franklin, J. (2010) Species traits affect the performance of species distribution models for plants in southern California. *Journal of Vegetation Science*, **21**, 177–189.
- Syphard, A.D., Radeloff, V.C., Keeley, J.E., Hawbaker, T.J., Clayton, M.K., Stewart, S.I. & Hammer, R.B. (2007) Human influence on California fire regimes. *Ecological Applications*, **17**, 1388–1402.
- Talmon, Y., Sternberg, M. & Grünzweig, J.M. (2011) Impact of rainfall manipulations and biotic controls on soil respiration in Mediterranean and desert ecosystems along an aridity gradient. *Global Change Biology*, **17**, 1108–1118.
- Valladares, F., Benavides, R., Rabasa, S., Pausas, J.G., Paula, S. & Díaz, M. (2013) Global change and Mediterranean forests: current impacts and potential responses. *Forests and global change* (ed. by D.A. Coomes, D.F.R.P. Burslem and W.D. Simonson), pp. 47–75. Cambridge University Press, Cambridge.
- Vargas, R., Sonnentag, O., Abramowitz, G., Carrara, A., Chen, J.M., Ciais, P., Correia, A., Keenan, T.F., Kobayashi, H., Ourcival, J.-M., Papale, D., Pearson, D., Pereira, J., Piao, S., Rambal, S. & Baldocchi, D. (2013) Drought influences the accuracy of simulated ecosystem fluxes: a model-data meta-analysis for Mediterranean oak woodlands. *Ecosystems*, **16**, 749–764.
- Vayreda, J., Martínez-Vilalta, J., Gracia, M. & Retana, J. (2012) Recent climate changes interact with stand structure and management to determine changes in tree carbon stocks in Spanish forests. *Global Change Biology*, **18**, 1028–1041.
- Verburg, P.H., van Berkel, D.B., van Doorn, A.M., van Eupen, M. & van den Heiligenberg, H.A. (2010) Trajectories of land use

- change in Europe: a model base exploration of rural futures. *Landscape Ecology*, **25**, 217–232.
- Viers, J.H., Fremier, A.K., Hutchinson, R.A., Quinn, J.F., Thorne, J.H. & Vaghti, M.G. (2012) Multiscale patterns of riparian plant diversity and implications for restoration. *Restoration Ecology*, **20**, 160–169.
- Vilà, M., Lloret, F., Ogheri, E. & Terradas, J. (2001) Positive fire–grass feedback in Mediterranean Basin woodlands. *Forest Ecology and Management*, **147**, 3–14.
- Vilà, M., Tessier, M., Suehs, C.M., Brundu, G., Carta, L., Galanidis, A., Lambdon, P., Manca, M., Médail, F., Moragues, E., Traveset, A., Troumbis, A.Y. & Hulme, P.E. (2006) Local and regional assessments of the impacts of plant invaders on vegetation structure and soil properties of Mediterranean islands. *Journal of Biogeography*, **33**, 853–861.
- Vilà-Cabrera, A., Martínez-Vilalta, J., Vayreda, J. & Retana, J. (2011) Structural and climatic determinants of demographic rates of Scots pine forests across the Iberian Peninsula. *Ecological Applications*, **21**, 1162–1172.
- Wain, A., Mills, G., McCaw, L. & Brown, T. (2008) Managing smoke from wildfires and prescribed burning in southern Australia. *Developments in environmental science* (ed. by A. Bytnerowicz, M.J. Arbaugh, A.R. Riebau and C. Andersen), pp. 535–550. Elsevier, Amsterdam.
- Westerling, A.L., Hidalgo, H.G., Cayan, D.R. & Swetnam, T.W. (2006) Warming and earlier spring increase western U.S. forest wildfire activity. *Science*, **313**, 940–943.
- Xu, L. & Baldocchi, D.D. (2004) Seasonal variation in carbon dioxide exchange over a Mediterranean annual grassland in California. *Agricultural and Forest Meteorology*, **123**, 79–96.
- Zaehle, S., Bondeau, A., Carter, T., Cramer, W., Erhard, M., Prentice, I.C., Reginster, I., Rounsevell, M.D.A., Sitch, S., Smith, B., Smith, P.C. & Sykes, M. (2007) Projected changes in terrestrial carbon storage in Europe under climate and land-use change, 1990–2100. *Ecosystems*, **10**, 380–401.
- Zampieri, M. & Lionello, P. (2011) Anthropogenic land use causes summer cooling in Central Europe. *Climate Research*, **46**, 255–268.
- Zhao, F., Zhang, L., Zongxue, X. & Scott, D. (2010) Evaluation of methods for estimating the effects of vegetation change and climate variability on streamflow. *Water Resources Research*, **46**, W03505.

BIOSKETCH

E. Doblas-Miranda is the research coordinator of the MONTES-Consolider project (<http://www.creaf.uab.es/montes/>), which aims to examine the relationship of mediterranean woodlands with the components of global change, and to identify opportunities to modify these components through appropriate woodland management. The present paper synthesizes most of the work carried out preparing the background of that project and analysing its outputs.

E.D.-M., J.R., J.M.-V. and P.L. conceived the ideas and principal scheme of the manuscript. E.D.-M. led the writing, with all authors contributing their different areas of expertise.

Editor: Martin Sykes